

BIOSECURITY NEW ZEALAND
IMPORTING COUNTRIES PHYTOSANITARY
REQUIREMENTS
REPUBLIC OF SOUTH AFRICA

Status: Approved

Date: 20 April 1999

Amendment record

Amendment No.	Date:	Nature of amendment:	Approved by:
1.	20 April 1999	Ammendment to sections 2.1, 2.6 and 4.1-4.6 inclusive. Re-issue of EPS.	SCO
2.	26 July1999	Change of address Section 1.1, addition of Section 4.1.2 (Frozen Fruit & Vegetables) and change of headings of Sections 4.6 and 4.6.1 (Herbs.... including Seed for Consumption).	SMN
3.	14 February 2003	Renaming and reformatting of standard. Amendment to Section 2.5 re MPLs.	WJH
4.	1 February 2005	Amendment of MAF contact details Section 1.1 and 1.2. Minor reformatting of document.	WJH

DISCLAIMER

The information in this standard is provided on the following basis. The phytosanitary requirements found in this standard are current at the time of publication and may be used as the basis of export certification. However, the requirements may be changed by importing countries at any time. MAF will make all reasonable efforts to advise exporters of these changes. However, MAF does not accept any liability for losses arising from importing countries failing to allow entry of plants or plant products due to changes in phytosanitary requirements of which MAF was either not advised or was not advised of within a reasonable period of time before their activation.

TABLE OF CONTENTS

1	GENERAL INFORMATION.....	3
1.1	FOR ENQUIRIES ABOUT THIS STANDARD CONTACT	ERROR! BOOKMARK NOT DEFINED.
2.	GENERAL REQUIREMENTS	5
2.1	IMPORT PERMITS	5
2.2	PHYTOSANITARY CERTIFICATES	5
2.3	QUARANTINE PESTS.....	6
2.4	INSPECTION ON ARRIVAL.....	14
2.5	MAXIMUM PEST LIMITS (MPL's).....	14
2.6	PROHIBITIONS.....	14
2.7	PORTS OF ENTRY.....	16
2.8	TRANSIT PROVISIONS.....	17
3.	COMMODITY CLASS REQUIREMENTS	17
4.	COMMODITY SPECIFIC REQUIREMENTS.....	17
4.1	FRUIT AND VEGETABLES	18
4.2	CUT FLOWERS/FOLIAGE.....	21
4.3	MISCELLANEOUS, PACKING MATERIAL AND PLANT (VEGETABLE) FIBRE.....	22
4.4	NURSERY STOCK	24
4.5	SEEDS (GRAIN)/NUTS	88
4.6	HERBS, SPICES, MEDICINAL PLANTS, & PLANTS FOR ANIMAL & HUMAN CONSUMPTION (INCLUDING SEED FOR CONSUMPTION)	115
4.7	WOOD, AND UNMANUFACTURED WOOD PRODUCTS	125

1 General Information

1.1 For enquires about this standard, contact:

Attn: Wayne Hartley
Exports Group
Biosecurity New Zealand
PO Box 2526
Wellington

Telephone: 64 4 498 9947
Facsimile: 64 4 498 9888
E-mail: wayne.hartley@maf.govt.nz

1.2 Phytosanitary Legislation

South Africa's phytosanitary requirements are documented in the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992) and is the document on which this export phytosanitary standard is based.

1.3 Definitions

The South Africa Agricultural Pests Act, 1983 (Act No. 36 of 1983) defines the following terms:

area of production any plot or group of plots that are cultivated as a unit and on which controlled goods are produced;

certificate the original specimen of a phytosanitary certificate or phytosanitary certificate of re-export that -

- (i) is in the form of the FAO model certificate;
- (ii) has been issued by the recognised authority in the country of production (ie. MAF) or the country of re-export of the controlled goods to which the certificate relates;

Controlled goods (a) any plant, pathogen, insect, exotic animal, growth medium, infectious thing, honey, beeswax or used apiary equipment;

(b) anything determined by the Minister in notice in the Republic of South Africa Government Gazette:

Growth medium any solid or liquid substance in which or on which plants are or can be cultivated, including soil in containers or structures used for cultivation of plants;

Import to introduce controlled goods into the Republic of South Africa or bring about the introduction of such goods into

the Republic of South Africa;

Infectious thing	anything except a plant, which may serve as a medium for the importation or spreading of any pathogen, insect or exotic animal;
Insect	any invertebrate member of the animal kingdom irrespective of the stage of development thereof but not such a member - (a) included in the definition of "fish" in section 1 of the Sea Fishery Act, 1988; or (b) that can affect man or an animal only (Definition of "insect" substituted by s. 1 (e) of Act 18 of 1989);
Officer	an officer or employee as defined in section 1 of the Public Service Act, 1984 (Act No. 111 of 1984);
Pathogen	any algae, bacterium, fungus, mycoplasma, rickettsia-like organism, spiroplasma, viroid, or virus, but does not include any such pathogen that can cause a disease in man or an animal only (Definition of "pathogen" substituted by s. 1 (h) of Act 18 of 1989);
Permit	a permit mentioned in section 3 (1) or 6 (3) of the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992);
Plant	any live or dead part of a plant and any derivatives of a plant;
Quarantine area	an area which has been determined as a quarantine area in an order under section 7 of the Agricultural Pests Act, 1983 (Act No. 36 of 1983) as amended by Agricultural Pests Amendment Act, 1992 (No. 9 of 1992);

2. General Requirements

2.1 Import Permits

Import permits are not required for the commodities specified in section 4.0 of this document unless it is specifically stated. All other commodities require an import permit. Where an import permit is required, phytosanitary information for controlled goods will be specified on the import permit.

2.2 Phytosanitary Certificates

Phytosanitary certificates are required for all commodities unless otherwise specified in section 4.0. Additional declarations and treatments for controlled goods which do not require import permits are specified in section 4.0 (Commodity Class information). For commodities requiring import permits, the phytosanitary requirements are specified on the import permit. Additional

declarations and treatments are to be entered in the relevant sections of the phytosanitary certificate.

2.3 Quarantine Pests

The following list of pathogens and pests are prohibited from entering the Republic of South Africa (refer also to the list of prohibited plants and invader plants):-

1. Abaca mosaic virus
2. Abutilon mosaic virus
3. *Acalitus essigi*
4. *Acalitus orthomera*
5. *Acarapis woodi*
6. *Aceria biopsida*
7. *Aceria breakeyi*
8. *Accria peucedamum*
9. *Aceria tulipae*
10. *Actinomyces ipomoeae*
11. *Aecidium fragiforme*
12. Alfalfa dwarf mosaic virus
13. *Alternaria cichorii*
14. *Alternaria padwickii*
15. *Anarsia lineatella*
16. *Anastrepha fraterculus*
17. *Anastrepha ludens*
18. *Anastrepha mombinpraeoptans*
19. *Anguina* spp.
20. *Anguina tritici*
21. Annual ryegrass toxicity syndrome
22. *Aphelenchoides besseyi*
23. *Aphelenchoides fragariae*
24. *Aphelenchoides ritzema-bosi*
25. Arabis mosaic virus
26. Artichoke curly dwarf virus
27. Artichoke mosaic virus
28. Artichoke mottle crinkle virus
29. *Ascochyta abelmoschi*
30. *Ascochyta gossypii*
31. *Ascochyta pisi*
32. *Ascochyta rabiei*
33. *Ascochyta* spp.
34. *Ascospaera apis*
35. Asparagus virus
36. *Aspergillus niger* f.sp. *floridanus*
37. *Aspidiella hartii*
38. Aster yellows mycoplasma
39. *Bacillus larvae*
40. Bacterial pathogens
41. *Balansia oryzae*
42. Banana bunchy top virus
43. Bean yellow Mosaic virus
44. Beet curly top virus

45. Black pepper stunt virus
46. Blueberry stunt virus
47. *Botrytis tulipae*
48. Broadbean stain virus
49. *Bursaphelenchus xylophilus*
50. *Cacoecimorpha pronubana*
51. Cadang-cadang viroid
52. *Caeoma sanctae-crucis*
53. *Caeoma torreya*
54. *Calacarus carinatus*
55. *Campanotus herculeanus*
56. Carnation etched ring virus
57. Carnation necrotic fleck virus
58. Carnation streak virus
59. *Cecidophyopsis ribis*
60. Celery mosaic virus
61. *Cephaleuros parasiticus*
62. *Cephalosporium diospyri*
63. *Cephalosporium maydis*
64. *Cephalosporium sacchari*
65. *Ceratocystis coerulescens*
66. *Ceratocystis fagacearum*
67. *Ceratocystis fimbriata*
68. *Ceratocystis fimbriata* f.sp. *platani*
69. *Ceratocystis* spp.
70. *Ceratocystis ulmi*
71. *Ceratostomella fimbriata*
72. *Cercospora angolensis* (Syn. *Phaeoramularia angolensis*)
73. *Cercospora coffeicola*
74. *Cercospora corchori*
75. *Cercospora ixorae*
76. *Cercospora mangifera*
77. *Cercospora pini-densiflorae*
78. *Cercospora purpurea*
79. *Cercospora sequioae*
80. *Cercospora sojina*
81. *Cercospora* spp.
82. *Cercospora transversiana*
83. *Ceutospora litchii*
84. *Chaetocnema pulicaria*
85. Cherry leafroll virus
86. Cherry rasp leaf virus
87. Chrysanthemum stunt viroid
88. *Chrysomyxa rhododendri*
89. *Ciboria betulae*
90. Citrus stubborn mycoplasma
91. *Claviceps purpurea*
92. *Claviceps* spp.
93. *Clitocybe tabescens*
94. Clostero viruses
95. Clover phyllody mycoplasma
96. *Cochliobolus miyabeanus* (Syn. *Helminthosporium oryzae*)
97. *Colletotrichum cajani*
98. *Colletotrichum capsici*

99. *Colletotrichum coffeanum* (CBD Strain)
100. *Colletotrichum dematium* f.sp. *truncatum*
101. *Colletotrichum fuscum*
102. *Colletotrichum lini*
103. *Colletotrichum panicicola*
104. *Colletotrichum villosum*
105. *Coniella diplodiella*
106. *Corynebacterium flaccumfaciens* pv. *betae*
107. *Corynebacterium flaccumfaciens* pv. *flaccumfaciens*
108. *Corynebacterium michiganense* pv. *michiganense*
109. *Corynebacterium michiganense* pv. *sepedonicum*
110. *Corynespora asiicola*
111. Cotton leaf curl virus
112. Cranberry false blossom mycoplasma
113. *Crinipellis palmivora*
114. *Crinipellis pernicioso*
115. *Cronartium coleosporioides*
116. *Cronartium comptoriae*
117. *Cronartium fusiforme*
118. *Cronartium quercuum*
119. *Cronartium strobultinum*
120. *Cryptospora longispora*
121. *Cryptostictis cupressi*
122. Cucumber leafroll virus
123. Cucumber mosaic virus
124. *Cydia molesta*
125. *Dacus cucurbitae*
126. *Dacus dorsalis*
127. *Dacus tryoni*
128. *Dacus zonatus*
129. *Diaporthe phaseolorum* f.sp. *batatatis*
130. *Diaporthe phaseolorum* f.sp. *caulivora*
131. *Didymella chrysanthemi*
132. *Ditylenchus destructor*
133. *Ditylenchus dipsaci*
134. *Ditylenchus* spp.
135. *Dothistroma pini*
136. *Dotichiza populea*
137. *Drechslera iridis*
138. *Drechslera maydis*
139. *Dysmicoccus brevipes*
140. *Echinodentium taxodii*
141. Elm mosaic virus
142. *Endothia gyrosa*
143. *Endothia parasitica*
144. *Ennomos subsignarius*
145. *Eotetranychus pruni* (*E. pomi*)
146. *Eotetranychus sexmaculatus*
147. *Ephelis pallida*
148. *Eriophyes gastronichus*
149. *Erwinia amylovora*
150. *Erwinia carotovora* pv. *musae*
151. *Erwinia carotovora* pv. *atroseptica*
152. *Erwinia chrysanthemi*

153. *Erwinia chrysanthemi* pv. *dianthi*
154. *Erwinia nigrifluens*
155. *Erwinia rhapontici*
156. *Erwinia rubrifaciens*
157. *Erwinia stewartii*
158. Euonymus variegation virus
159. Euphorbia mosaic virus
160. *Exobasidium camelliae*
161. *Exobasidium vexans*
162. *Frankliniella occidentalis*
163. Fungal pathogens
164. *Fusarium oxysporum* f.sp. *cubense*
165. *Fusarium oxysporum* f. sp. *lycopersici* Race III
166. *Fusarium oxysporum* f. sp. *psidii*
167. *Fusarium oxysporum* f.sp. *zingiberi*
168. *Ganoderma lucidum*
169. *Globodera* spp.
170. *Gloeotinia temulenta*
171. *Glomerella cingulata*
172. *Gnomonia platani* (syn *G. veneta*)
173. *Godrona cassandrae*
174. *Goplana dioscorea*
175. *Gremmeniella abeitina*
176. Guar top necrosis virus
177. *Guignardia aesculi*
178. *Guignardia bidwelii*
179. *Gymnosporangium* spp.
180. *Hendersonia agathi*
181. *Heterodera glycines*
182. *Heterodera humuli*
183. *Heterodera* spp.
184. Hop mosaic virus
185. Hop nettle virus complex
186. Hydrangea ringspot virus
187. *Hymenochaeta mongeotii*
188. *Hymenoptera* spp.
189. Iilar group viruses
190. *Kabathina thujae*
191. *Kabatiella nigricans*
192. *Kabatiella zeae*
193. *Keithia thujina*
194. *Koleroga noxia*
195. Leaf scorch
196. *Leptinotarsa decemlineata*
197. *Leptosphaeria maculans* (Syn. *Phoma lingam*)
198. *Leptosphaeria obtusispora*
199. Lethal yellowing mycoplasma
200. Lettuce mosaic virus
201. Lily mosaic virus
202. Lily rosette virus
203. *Liriomyza* spp.
204. *Lophodermium cedri*
205. *Lophodermium* spp.
206. *Lumbricus rubellus*

207. *Lymantria disparina*
208. *Magnaporthe salvinii*
209. *Malpigamoeba mellificae*
210. *Mampava rhodoneura*
211. *Marssonina brannea*
212. *Melampsora hypericorum*
213. *Melampsora pinitorqua*
214. *Meloidogyne* spp.
215. *Microcyclus ulei*
216. *Microsphaeria alni*
217. Mollusca
218. *Monilia roreri*
219. *Monilochaetes infuscans*
220. Mosaic
221. Mosaic virus infection
222. *Mycosphaerella citri*
223. *Mycosphaerella fijiensis*
224. *Mycosphaerella linorum*
225. Myrobolan latent ringspot virus
226. Narcissus yellow stripe virus
227. Necrotic ringspot virus serotypes
228. *Nectria cinnabarina*
229. *Nectria galligena*
230. Nematodes (plant parasitic/plant parasites)
231. *Neovossia indica*
232. *Oidium euonymi-japonici*
233. *Opogona sacchari*
234. *Ovulina azaleae*
235. *Paramyelois transitella*
236. *Pardalaspis cyanescens*
237. Pea seedborne mosaic virus
238. Peanut marginal chlorosis virus
239. Peanut mottle virus (syn. Groundnut mottle virus)
240. Peanut stunt virus
241. Pelargonium leaf curl virus
242. Peperomia ringspot virus
243. *Peronosclerospora maydis*
244. *Peronosclerospora phillipinensis*
245. *Peronosclerospora sacchari*
246. *Peronosclerospora spontanea*
247. *Peridermium cedri*
248. *Peronospora arborescens*
249. *Peronospora documeti*
250. *Peronospora farinosa*
251. *Peronospora hyoschyami*
252. *Peronospora jacksonii*
253. *Peronospora mesembryanthemi*
254. *Peronospora schachtii*
255. *Peronospora tabacina*
256. *Phakopsora cheoana*
257. *Phakopsora jatrophiicola*
258. *Phakopsora pachyrhizi*
259. *Phakopsora zizyphi-vulgaris*
260. *Phoma cisti*

261. *Phoma sabdariffae*
262. *Phoma* spp.
263. *Phoma strasseri*
264. *Phomopsis annonacearum*
265. *Phomopsis heveae*
266. *Phomopsis theae*
267. *Phomopsis vexans*
268. Phormium yellowing mycoplasma
269. *Phyllosticta dracaenae*
270. *Phymatotrichum omnivorum*
271. *Physalospora miyabeana*
272. *Physopella ampelopsidis*
273. Phytophagous mites
274. *Phytophthora capsici*
275. *Phytophthora cryptogea*
276. *Phytophthora fragariae*
277. *Phytophthora infestans*
278. *Phytophthora palmivora*
279. *Phytophthora* spp.
280. Plant parasitic nematodes
281. *Plasmodiophora brassicae*
282. *Plasmopara chrysanthemi-coronarii*
283. *Plasmopara halstedii*
284. *Pleospora papaveracea*
285. *Ploioderma lethale*
286. Plum Pox virus (Sharka)
287. Poinsettia mosaic virus
288. *Polaccia saliciperda*
289. *Polyporus gilvus*
290. *Polyporus sanguineus*
291. *Polyscytalum pustulans*
292. Potato spindle tuber viroid
293. *Pratylenchus brachyurus*
294. *Pratylenchus scribneri*
295. *Prostephanus truncatus*
296. *Pseudococcus citri*
297. *Pseudomonas aleuritides*
298. *Pseudomonas andropogonis*
299. *Pseudomonas aptata*
300. *Pseudomonas caryophylli*
301. *Pseudomonas maublanci*
302. *Pseudomonas phaseolicola*
303. *Pseudomonas saliciperda*
304. *Pseudomonas solanacearum*
305. *Pseudomonas solanacearum* biotypes III and IV
306. *Pseudomonas syringae* pv. *mellea*
307. *Pseudomonas syringae* pv. *lisi*
308. *Pseudomonas syringae* pv. *populae*
309. *Pseudomonas syringae* pv. *savastanoi*
310. *Pseudomonas viridiflava*
311. *Pseudoperonospora humuli*
312. *Psylla pyricola*
313. *Puccinia cari-bistortea*
314. *Puccinia graminis*

315. *Puccinia horiana*
316. *Puccinia psidii*
317. *Puccinia* spp.
318. *Pucciniastrum actinidiae*
319. *Pyrenochaeta phlogina*
320. *Pythium myriotylum*
321. *Radopholus citrophilis*
322. *Radopholus similis*
323. *Ramularia bellulensis*
324. *Ramularia cyclaminicola*
325. *Rhadinaphelenchus cocophilus*
326. *Rhagoletis cerasi*
327. *Rhagoletis pomonella*
328. *Rhizoctonia solani*
329. *Rhyncophorus palmarum*
330. Root wilt
331. Rose Rosette virus
332. Rose wilt virus
333. *Rosselinia bunodes*
334. *Rotylenchus reniformis*
335. *Sanninoidea exitiosa*
336. *Sclerophthora rayssiae* f.sp. *zeae*
337. *Sclerotinia bulborum*
338. *Sclerotinia convulata*
339. *Sclerotinia narcissicola*
340. *Sclerotinia polyblastis*
341. *Sclerotinia ricini*
342. *Sclerotinia trifolium*
343. *Sclerotium tuliparum*
344. *Seiridium cardinale*
345. *Septobasidium aleuritides*
346. *Septoria azaleae*
347. *Septoria gladioli*
348. *Septoria glycines*
349. Shuck die-back disease
350. *Sirex noctilio*
351. Soil-borne viruses
352. *Sorosporium syntherismae*
353. Soybean mild mosaic virus
354. Soybean stunt virus
355. *Sphaceloma manihoticola*
356. *Sphaceloma punicae*
357. *Sphaerotheca mors-uvae*
358. *Sphaerulina taxicola*
359. *Spongospora* spp.
360. *Steneotarsonemus laticeps*
361. Strawberry latent ringspot virus
362. *Streptomyces scabies*
363. Sugarcane Fiji virus
364. Sugarcane grassy shoot virus
365. Sugarcane sereh virus
366. *Synchytrium endobioticum*
367. *Synchytrium piperi*
368. *Tetranychus canadensis*

369. *Tetranychus mcdanielli*
370. *Tetranychus pacificus*
371. *Tetranychus schoenei*
372. *Tetranychus viennensis*
373. *Thyronectria denigrata*
374. *Tilletia barclayana*
375. *Tilletia contraversa*
376. Tobacco leaf curl virus
377. Tobacco rattle virus
378. Tobacco ringspot virus
379. Tobacco streak virus
380. Tomato aspermy virus
381. Tomato black ring virus
382. Tomato ringspot virus
383. *Trioza calacarus*
384. *Trogoderma granarium*
385. *Tropilaelaps clareae*
386. Tulip white streak virus
387. Uredinales
388. Uredinales (Pine rusts)
389. *Uredo phormii*
390. *Urocystis agropyri*
391. *Urocystis cepulae* (Syn. *U. colchici*)
392. *Uromyces cytisi*
393. *Uromyces genistae-tinctoriae*
394. *Uromyces* spp.
395. *Ustilaginoidea virens*
396. *Ustilago coicis*
397. *Ustilago nuda*
398. *Ustilago utriculosa*
399. *Ustilago violacea*
400. *Varroa jacobsoni*
401. *Verticillium albo-atrum*
402. *Verticillium dahliae*
403. *Verticillium* spp.
404. Virus and virus-like diseases affecting honey bees
405. Virus and virus diseases
406. Virus chlorosis
407. Wilt diseases caused by fungi
408. *Xanthomonas campestris* pv. *begoniae*
409. *Xanthomonas campestris* pv. *campestris*
410. *Xanthomonas campestris* pv. *cassavae*
411. *Xanthomonas campestris* pv. *cassiae*
412. *Xanthomonas campestris* pv. *celebensis*
413. *Xanthomonas campestris* pv. *citri*
414. *Xanthomonas campestris* pv. *corylina*
415. *Xanthomonas campestris* pv. *erythrinae*
416. *Xanthomonas campestris* pv. *hyacinthi*
417. *Xanthomonas campestris* pv. *khayae*
418. *Xanthomonas campestris* pv. *manihotis*
419. *Xanthomonas campestris* pv. *oryzae*
420. *Xanthomonas campestris* pv. *oryzicola*
421. *Xanthomonas campestris* pv. *phormicola*
422. *Xanthomonas campestris* pv. *vasculorum*

- 423. *Xanthomonas campestris* pv. *vesicatoria*
- 424. *Xanthomonas fragariae*
- 425. *Xanthomonas gorlincovianum*
- 426. *Xanthomonas* spp.
- 427. *Xanthomonas panici*
- 428. Yam mosaic virus

2.4 Inspection on Arrival

All commodities are subject to inspection on arrival. The level of inspection conducted by South Africa is not known by MAF.

2.5 Maximum Pest Limits (MPL's)

For all commodities exported to South Africa requiring phytosanitary certificates, MPL's are:

Quarantine pests* specified by South Africa	0.5%
	Soil 25g/600unit

*Quarantine pests for South Africa include organisms identified within:

- Section 2.4 of this standard
- Additional declarations
- Phytosanitary import permit

2.6 Prohibitions

Any live plant, vegetative propagative material, or a seed of a kind of plant that has been declared a weed or invader plant is prohibited entry to the Republic of South Africa except by special import permit issued only to a person in charge of a *bona fide* research institution (Agricultural Pets Act 1983 (Act 36 of 1983). Soil (all types) and sand is prohibited from entering the Republic of South Africa.

Declared Weeds include:

Albizia lophantha
Alhagi camelorum
Caesalpinia decapetala
Cannabis sativa
Cereus peruvianus
Cestrum aurantiacum
Cestrum laevigatum
Cestrum parqui
Chromolaena odorata
Cirsium vulgare
Cuscuta campestris
Datura ferox
Datura stramonium
Elodea spp. and other plants of the family Hydrocharitaceae
Eichhornia spp. and other plants of the family Pontederiaceae
Hakea gibbosa

Hakea sericea
Hakea suaveolens
Harrisia martinii
Lantana camara and any entity which have partly or wholly been derived from the *Lantana camara* complex by means of hybridisation or selection under natural or artificial conditions.
Lemna spp. and other plants of the family Lemnaceae
Myriophyllum spp. and other plants of the family Haloragidaceae
Opuntia aurantiaca
Opuntia dillenii
Opuntia exaltata
Opuntia ficus-indica excluding all spineless cultivars and selections
Opuntia imbricata
Opuntia lindheimeri
Opuntia rosea
Opuntia spinulifera
Opuntia stricta
Opuntia vulgaris
Orobanche minor
Pereskia aculeata
Pereskia grandifolia
Pereskia stratiotes
Rubus cuneifolius
Salvinia spp. and other plants of the family Salviniaceae
Sesbania punicea
Solanum elaeagnifolium
Solanum mauritianum
Solanum sisymbriifolium
Stipa tenuissima
Stipa trichotoma
Xanthium spinosum
Xanthium strumarium

Declared Invader Plants include:-

Acacia caffra
Acacia cyclops.
Acacia dealbata
Acacia erubescens
Acacia fleckii
Acacia hebeclada
Acacia karroo
Acacia longifolia
Acacia mearnsii
Acacia melanoxylon
Acacia mellifera
Acacia nigrescens
Acacia nilotica
Acacia reficiens subsp. *reficiens*
Acacia robusta subsp. *robusta*
Acacia saligna
Acacia senegal var. *rostrata*
Acacia tenuispina
Acacia tortilis subsp. *heteracantha*
Colophospermum mopane

Combretum apiculatum subsp. *apiculatum*
Commiphora pyracanthoides subsp. *pyracanthoides*
Dichrostachys cinerea subsp. *africana*
Grewia bicolor
Grewia flava
Grewia flavescens
Leptospermum laevigatum
Maytenus senegalensis
Ochna pulchra
Pinus pinaster
Prosopis spp.
Rhigozum trichotomum
Rhus lancea
Tarchonanthus camphoratus
Terminalia sericea

2.7 Ports of Entry

A person importing controlled goods into the Republic of South Africa on the authority of a permit shall do so only through a prescribed port of entry, except where an executive officer has determined some other place.

Prescribed ports include:-

The harbours of :-

Cape Town
Durban
East London
Port Elizabeth

The container depots of :-

Cape Town
City Deep
Durban
Port Elizabeth

The airports of :-

D. F. Malan (Cape Town)
H. F. Verwoerd (Port Elizabeth)
Jan Smuts (Johannesburg)
Lois Botha (Durban)

The border control ports of :-

Vioolsdrif
Nakop
Grobler's Bridge
Kopfontein Gate
Schilpad's Gate
Ramatlabama
Beit Bridge
Lebombo
Border Gate (Managa)
Oshoek
Nerston
Mahamba
Golela
Ficksburg Bridge
Maseru Bridge

The Main Post Offices at:-

Cape Town
Durban
Johannesburg
East London
Port Elizabeth
Pretoria

2.8 Transit Provisions

The Republic of South Africa have not stipulated transit details for controlled goods.

3. Commodity Class Requirements

Import permits are not required for the commodities specified in section 4.0 unless it is otherwise specified. All other commodities require an import permit. Where an import permit is required phytosanitary information for controlled goods will be specified on the import permit. Phytosanitary certificates are required for all commodities unless otherwise specified. Additional declarations and treatments for controlled goods which do not require import permits are specified in section 4.0 (Commodity Class information). Additional declarations and treatments are to be entered in the relevant sections of the phytosanitary certificate.

4. Commodity Specific Requirements

This section has been split into commodity classes, and further divided into commodity subclasses where necessary. Within each commodity class relevant information on general requirements for import permits (section 2.1), phytosanitary certificates (section 2.2), inspection (section 2.3), prohibitions (section 2.5) and ports of entry (section 2.6) should be referred to with commodity class specific phytosanitary information.

4.1 Fruit and Vegetables

4.1.1 Dried/fresh fruit and vegetables

Does not include seed for consumption or processing.
Phytosanitary import permit required.

4.1.2 Frozen fruit and vegetables

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Frozen fruit and vegetables require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Actinidia</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , & <i>Dacus dorsalis</i> . <i>Cydia molesta</i> does not occur on kiwifruit in New Zealand. (Please note: This AD to be verified by MAF).
<i>Ananas</i>	AD3:- The area of production is free from <i>Dysmicoccus brevipes</i> AD4: - The consignment is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cydia molesta</i> , & <i>Dacus dorsalis</i> .
<i>Artocarpus</i>	AD2: New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Asimina</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Averrhoa</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Bambusa</i>	AD11: - The consignment is free from soil or growth mediums. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Capsicum</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Carica</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> & <i>Dacus tryoni</i> .

<i>Citrus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cercospora angolensis</i> (Syn. <i>Phaeoramularia angolensis</i>), <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> , <i>Mycosphaerella citri</i> & <i>Xanthomonas campestris pv. citri</i> .
<i>Citrus</i> (for Jewish festivals)	No additional declarations required. T8 = the consignment was treated by immersion in sodium hypochlorite. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Citrus</i> (Candied peel)	AD2: - New Zealand is free from <i>Cercospora angolensis</i> (Syn. <i>Phaeoramularia angolensis</i>), and <i>Xanthomonas campestris pv. citri</i> .
<i>Cocos</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Colchicum</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
Cucurbitaceae	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> & <i>Dacus tryoni</i> .
<i>Curcuma</i>	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Dioscorea</i>	AD11: - The consignment is free from soil or growth mediums. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Diospyros</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> .
<i>Fragaria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .
<i>Litchi</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .
<i>Lycopersicon</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , & <i>Pardalaspis cyanescens</i> .
<i>Malus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , & <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> , <i>Eotetranychus sexmaculatus</i> & <i>Rhagoletis pomonella</i> .

<i>Manilkara</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Myricaria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Olea</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Passiflora</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Persea</i>	AD2:- New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Phaseolus</i>	Phytosanitary certificate required.
<i>Phoenix</i> (with seed)	No additional declarations required. (The importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation).
<i>Phoenix</i> (stoned)	Phytosanitary certificate required.
<i>Physalis</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Pisum</i>	Phytosanitary certificate required.
<i>Pouteria</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> .
<i>Psidium</i>	AD2: - New Zealand is free from <i>Aceria biopsida</i> , <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> & <i>Eotetranychus sexmaculatus</i> . AD3:- The area of production is free from <i>Puccinia psidii</i> . AD4: - The consignment is free from <i>Cydia molesta</i> .
<i>Punica</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Pyrus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
<i>Ribes</i>	AD2: - New Zealand is free from <i>Aceria breakeyi</i> , <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Cecidophyopsis ribis</i> , <i>Dacus dorsalis</i> , & <i>Dacus tryoni</i> .

Rosaceae (Genera & sp. not mentioned elsewhere)	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
<i>Rubus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Mycosphaerella citri</i> . AD4: - The consignment is free from <i>Cydia molesta</i> and <i>Psylla pyricola</i> .
Rutaceae	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Mycosphaerella citri</i> & <i>Xanthomonas campestris</i> pv. <i>citri</i> .
<i>Solanum</i> (excluding <i>S. tuberosum</i>)	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> and <i>Dacus tryoni</i> .
<i>Zea</i>	Phytosanitary certificate required.
<i>Zizyphus</i>	AD2: - New Zealand is free from <i>Anastrepha fraterculus</i> , <i>Anastrepha ludens</i> , <i>Anastrepha mombinpraeoptans</i> , <i>Dacus dorsalis</i> , <i>Dacus tryoni</i> , <i>Eotetranychus pruni</i> (<i>Eotetranychus pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Rhagoletis cerasi</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD3: - The area of production is free from <i>Erwinia amylovora</i> , and <i>Nectria galligena</i> . AD4: - The consignment is free from <i>Cydia molesta</i> .

4.2 Cut Flowers/Foliage

4.2.1 Dried/fresh cut flowers/foliage

Phytosanitary import permit required.

4.3 Miscellaneous, Packing Material and Plant (Vegetable) Fibre

4.3.1 Miscellaneous and packing material

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Miscellaneous and packing material require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

Compost, compost inoculum, vermi compost	AD:- Either "Laboratory examination of representative samples has shown the consignment to be free from plant pathogenic - bacteria, fungi, nematodes and soil." or "The consignment has been sterilised."
Bamboo/Rattan	No additional declarations required.
Bark (of <i>Barcena</i> spp. only)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
Cork (unmanufactured)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
Cork (manufactured)	No additional declarations required.
Growth media & Orchid rooting media	(excluding bark of coniferous plants).AD:- Either "Laboratory examination of representative samples has shown the consignment to be free from plant pathogenic - bacteria, fungi, nematodes and soil." or "The consignment has been sterilised."
Resins/Vegetable gums	No additional declarations required.
Sawdust	Import permit required.
Tobacco (leaf, stalks etc.)	Import permit required.

4.3.2 Plant (Vegetable) Fibre

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Plant (Vegetable) Fibre require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Bombacaceae</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. Must contain less than 1 seed per 500g fibre.
<i>Cocos</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).
<i>Corchoris</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).
<i>Gossypium</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. Must contain less than 1 seed per 2kg fibre.
<i>Hibiscus</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.
<i>Leopoldiana piassuba</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas (if necessary).
<i>Linum</i>	No additional declarations required.
<i>Sorghum</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas.

4.4 Nursery Stock

4.4.1 Bud wood/Cuttings

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Bud wood/Cuttings require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Abelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acalypha</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acanthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Achillea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Acokanthera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Actinidia</i>	AD4:- The consignment is free from <i>Eotetranychus sexmaculatus</i> , <i>Pseudomonas viridiflava</i> and <i>Pucciniastrum actinidiae</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Actinodium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Adenanthos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Adenia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Adenium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Agave</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , and <i>Steneotarsonemus laticeps</i> . T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aglaonema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.

<i>Ajuga</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Alberta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aleurites</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas aleuritides</i> , <i>Septobasidium aleuritides</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aloe</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Polyporus sanguineus</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Amaranthaceae	AD4:- The consignment is free from Beet curly top virus, Cucumber mosaic virus, Tobacco ringspot virus, Tomato black ring virus and Tomato ringspot virus.
<i>Amphicarpaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Androsace</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Angelica</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Anigozanthos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Anthurium</i>	AD4:- The consignment is free from <i>Radopholus citrophilis</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Antigonon</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aphelandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aquilegia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, & Cucumber mosaic virus did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Arabis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria maculans</i> (Syn. <i>Phoma lingam</i>) & <i>Xanthomonas campestris pv. campestris</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum fumigant.
<i>Aralia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Arbutus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ardisia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Arisaema</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Aristolochia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Armeria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Armoracia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Celery mosaic virus, <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Artemisia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Plasmopara chrysanthemi-coronarii</i> , & <i>Plasmopara halstedii</i> , did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Artocarpus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Koleroga noxia</i> & <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Asterosperma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Atropa</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora hyoschymi</i> , & Tobacco rattle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Azadirachta</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ganoderma lucidum</i> & <i>Polyporus gilvus</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Azalea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Baeckea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Banksia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Barcena</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bauera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bauhinia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Beaucarnea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Beaufortia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Belaperone</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bellis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus did not occur on those plants; or (ii) does not occur in the area of production concerned.
<i>Bischofia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bougainvillea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Bouvardia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Brachychiton</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Brighamia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Buddleia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calathea</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> .
<i>Callistemon</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Callitris</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Cercospora sequioae</i> , <i>Cryptostictis cupressi</i> , <i>Gymnosporangium</i> spp., <i>Kabathina thujae</i> , & <i>Seiridium cardinale</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Calothamnus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calpurnia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calycanthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Calytrix</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Camellia</i>	AD2:- The area of production is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , & <i>Exobasidium vexans</i> AD4:- The consignment is free from <i>Cephaleuros parasiticus</i> , <i>Phomopsis theae</i> , & <i>Xanthomonas gortencovianum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant. (The material is to be of elite stock.)
<i>Campanula</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from Uredinales (Pine rusts).
<i>Cananga</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Capparis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Caragana</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Carissa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cassiope</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cedrela</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora cheoana</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceratonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceratopetalum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ceropegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Chamaelaucium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cheiranthus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Plasmiodiophora brassicae</i> & <i>Xanthomonas campestris</i> pv. <i>campestris</i> did not occur on those plants; or (ii) do not occur in the area of production concerned.
<i>Cicer</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ascochyta rabiei</i> , Beet curly top virus, <i>Verticillium albo-atrum</i> , & <i>Xanthomonas campestris</i> pv. <i>cassiae</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Cistus</i>	AD4:- The consignment is free from <i>Phoma cisti</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Clematis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Clerodendrum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cneorum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Codonanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Coleus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cordyline</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> & <i>Phyllosticta dracaenae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Correa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cortaderia</i>	No additional declarations required. T6 = the consignment concerned was treated by an appropriate hot-water treatment.
<i>Costus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crescentia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crinodendron</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Crossandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ctenanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Cussonia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
Cycadales	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cytissus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Uromyces cytisi</i> , & <i>Uromyces genistae-tinctoriae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Dahlia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> , & fungal wilt diseases did not occur on those plants or (ii) do not occur in the area of production concerned.

<i>Dalbergia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Danae</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Daphne</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Dasyliiron</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Davidia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Delonix</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Delphinium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Beet curly top virus, & Tobacco ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Diascia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dieffenbachia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , <i>Phytophthora</i> spp. did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Dionaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dionysia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dioscorea</i>	AD4:- The consignment is free from <i>Goplana dioscorea</i> , & Yam mosaic virus.
<i>Dipladenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Doryanthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Draba</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Dracaena</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> , & <i>Phyllosticta dracaenae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Echeveria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Edgeworthia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Elaeagnus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria galligena</i> , & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.

<i>Epacris</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ephedra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Epipremnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Eremaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Erica</i>	AD4:- The consignment is free from <i>Phytophthora</i> spp., & <i>Verticillium</i> spp.
<i>Erigeron</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Eryngium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Accria peucedamum</i> , & <i>Aceria tulipae</i> .
<i>Erythrina</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Xanthomonas campestris</i> pv. <i>erythrinae</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Escallonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Euonymus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euonymus</i> variegation virus, & <i>Oidium euonymi-japonici</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Euphorbia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euphorbia</i> mosaic virus, & <i>Poinsettia</i> mosaic virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Euptelea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Eustoma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Exacum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Fittonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Flindersia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Hymenochaeta mongeotii</i> , did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Fremontia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Fuchsia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Gardenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Geraniaceae	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Pelargonium leaf curl virus, Tomato black ring virus, & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Gleditsia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Thyronectria denigrata</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Gmelina</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Grevillea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Halimocistus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hamamelis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> & <i>Polyporus gilvus</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Hebe</i>	T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hedera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Helianthemum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hemiandra</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide , T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hibbertia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hosta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hoya</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Hydrangea</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Hydrangea ringspot virus, Tobacco ringspot virus, & Tomato ringspot virus, did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hyoscyamus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora hyoschyami</i> , & Tobacco rattle virus did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hypoestes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Hyssopus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ilex</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Jatropha</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora jatrophiicola</i> , and <i>Phytophthora palmivora</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Kalanchoe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kalmia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Khaya</i>	AD4:- The consignment is free from <i>Xanthomonas campestris pv. khayae</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kolkwitzia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Arabis mosaic virus, Cherry leafroll virus, & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Kunzea</i>	AD:- T1 = treated with a wide spectrum fungicide for control of <i>Phakopsora pachyrhizi</i> .
<i>Laburnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Tobacco ringspot virus, & <i>Uromyces genistae-tinctoriae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Lambertia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Euphorbia mosaic virus, & Poinsettia mosaic virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Lamium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Laurelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Lavandula</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Leschenaultia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Limonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Liquidambar</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Liriodendron</i>	AD3:- The area of production is free from <i>Ceratocystis coerulescens</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lisianthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lonicera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Lychnis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lycopodium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lysimachia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Lythrum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Magnolia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned, and <i>Nectria galligena</i> does not occur within 1 km of the area of production.
<i>Malva</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Abutilon mosaic virus did not occur on those plants or (ii) does not occur in the area of production concerned. AD2:- :- New Zealand is free from Cotton leaf curl virus.
<i>Manilkara</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Medinilla</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Menianthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mesembryanthemum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora mesembryanthemi</i> did not occur or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Metrosideros</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mikania</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mimulus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora jacksonii</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Mimusops</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> , did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Mitella</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Monarda</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Moringa</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Monstera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Mussaenda</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Myrmecodia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nepenthes</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nepeta</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Nertera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Oenanthe</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Oenothera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phymatotrichum omnivorum</i> , & <i>Verticillium albo-atrum</i> , did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Olearia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Onobrychis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Origanum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Patersonia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia convulata</i> , <i>Sclerotium tuliparum</i> , & Tobacco rattle virus did not occur on those plants or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> & <i>Ditylenchus dipsaci</i> .
<i>Peltiphyllum</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> & <i>Synchytrium endobioticum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Penstemon</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus, and <i>Phymatotrichum omnivorum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Peperomia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus, and Peperomia ringspot virus did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Persoonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Petasites</i>	AD:- T1 = treated with a wide spectrum fungicide for control of <i>Puccinia</i> spp. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Philadelphus</i>	AD4:- The consignment is free from Uredinales. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Philodendron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Phlomis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Phlox</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pyrenochaeta phlogina</i> , Tomato aspermy virus, & <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Phormium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Phormium yellowing mycoplasma, <i>Uredo phormii</i> , & <i>Xanthomonas campestris</i> pv. <i>phormicola</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Phylloglossum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Physostegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Phyteuma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pimpinella</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Puccinia cari-bistortea</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD:- T1 = treated with a wide spectrum fungicide for control of <i>Puccinia cari-bistortea</i> . T2 = treated with a wide spectrum insecticide or fumigant.

<i>Pisonia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pittosporum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Platanus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ceratocystis fimbriata f.sp. platani</i> , & <i>Gnomonia platani</i> (syn. <i>G. veneta</i>), did not occur on those plants or (ii) do not occur in the area of production concerned. T2 = treated with a wide spectrum insecticide or fumigant.
<i>Plectranthus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Plumeria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polemonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polygonum</i>	AD4:- The consignment is free from <i>Uredo phormii</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
Polypodiaceae	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Polyscias</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Pothos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pouteria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Preslia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Prostanthera</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Prunella</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Psilotum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pulmonaria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Pyrostegia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rhamnus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria cinnabarina</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Rhododendron</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Rhodohypoxis</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rhoeo</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rivinia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Rosmarinus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ruellia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas solanacearum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Rumex</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Ruscus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Salix</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Physalospora miyabeana</i> , <i>Polaccia saliciperda</i> , <i>Pseudomonas saliciperda</i> , & Virus chlorosis did not occur on those plants or (ii) do not occur in the area of production concerned, and <i>Nectria galligena</i> did not occur within 1 Km of the place of production. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Salvia</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phymatotrichum omnivorum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sambucus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus, Cherry leafroll virus & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Sansevieria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sarcocephalus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sarracenia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Saxifraga</i>	AD4:- The consignment is free from Uredinales. AD:- T1 = treated with a wide spectrum fungicide for control of Uredinales.
<i>Scaevola</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Schefflera</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned.

<i>Schizaea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Scindapsus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Simmondsia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Soleirolia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Sonerila</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Spathiphyllum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stachys</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stapelia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Statice</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Stephanotis</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysanthemum</i> stunt viroid did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Stereospermum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Symphoricarpos</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Syngonium</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Tabebuia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Talinum</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Telopia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Thunbergia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tilia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tmesipterus</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tolmiea</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

<i>Trachelium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tradescantia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Tryphonium</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Veronica</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Verticordia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Viburnum</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants or (ii) does not occur in the area of production concerned. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Weigela</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus Cherry leafroll virus & Tomato ringspot virus did not occur on those plants or (ii) do not occur in the area of production concerned.
<i>Wisteria</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Wistringia</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Xanthosoma</i>	No additional declarations required. T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.
<i>Yucca</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria obtusifolia</i> did not occur on those plants or (ii) does not occur in the area of production concerned.
<i>Zizyphus</i>	ADI:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora zizyphi-vulgaris</i> did not occur on those plants or (ii) does not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> . T1 = treated with a wide spectrum fungicide, T2 = treated with a wide spectrum insecticide or fumigant.

4.4.2 Bulbs/Corms/Rhizomes/Tubers etc for propagation

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Bulbs/Corms/ Rhizomes /Tubers etc require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Aconitum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Agapanthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Aglaonema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Alstroemeria</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Amaryllidaceae (excluding species mentioned elsewhere)	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Pratylenchus scribneri</i> , <i>Sclerotium tuliparum</i> , & <i>Steneotarsonemus laticeps</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Androsace</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

- Aquilegia* AD3:- The area of production is free from Beet curly top virus, Cucumber mosaic virus, *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Asparagus* AD2:- New Zealand is free from Asparagus virus. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Astilbe* AD2:- New Zealand is free from Asparagus virus. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Begonia* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Xanthomonas campestris pv. begoniae* did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Beilis* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Caladium* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Sclerotinia bulborum*, & *Sclerotium tuliparum* did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Cephalotus* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Sclerotinia bulborum*, *Sclerotium tuliparum* did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor* & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.

<i>Clivia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> <i>Pratylenchus scribneri</i> <i>Sclerotium tuliparum</i> , <i>Steneotarsonemus laticeps</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Colchicum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotium tuliparum</i> , <i>Urocystis cepulae</i> (Syn. <i>U. colchici</i>) did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cyclamen</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ramularia cyclaminicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Cynara</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Artichoke curly dwarf virus, Artichoke mosaic virus, Artichoke mottle crinkle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> and <i>Rotylenchus reniformis</i> AD11:- The consignment is free from soil or growth mediums.
<i>Dahlia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> & fungal wilt diseases did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , and <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Darlingtonia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Dicentra</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Eryngium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Accria peucedanum</i> , <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , and <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Gloriosa</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Heliotropium</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Hyacinthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , <i>Sclerotium tuliparum</i> , & <i>Xanthomonas campestris pv. hyacinthi</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Incarvillea</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Iridaceae (excluding species mentioned elsewhere)	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Drechslera iridis</i> , <i>Sclerotinia bulborum</i> , <i>Sclerotium tuliparum</i> , & Tobacco rattle virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Lewisia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Liatriis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

Liliaceae (excluding species mentioned elsewhere)	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Lily mosaic virus, Lily rosette virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Macropidia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> , & <i>Sclerotium tuliparum</i> , did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from Lily mosaic virus, Lily rosette virus, <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
Marantaceae	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus, & <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Mirabilis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Mitella</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Myrmecodia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Oxalis</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

<i>Paeonia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Panax</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Colletotrichum panicicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Phlomis</i>	AD3:- The area of production is free from <i>Phomopsis vexans</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Phytax</i>	AD3:- The area of production is free from <i>Phomopsis vexans</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> & Uredinales. AD11:- The consignment is free from soil or growth mediums.
<i>Sandesonia</i>	Refer to requirements Liliaceae.
<i>Sarracenia</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Strelitzia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus, Banana bunchy top virus, <i>Erwinia carotovora</i> pv. <i>musae</i> , & <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Mycosphaerella fijiensis</i> , <i>Phymatotrichum omnivorum</i> , <i>Pseudomonas solanacearum</i> <i>Synchytrium endobioticum</i> , & <i>Xanthomonas campestris</i> pv. <i>celebensis</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Radopholus citrophilus</i> , & <i>Radopholus similis</i> . AD11:- The consignment is free from soil or growth mediums.
<i>Tacca</i>	AD3:- The area of production is free from <i>Phymatotrichum omnivorum</i> , & <i>Synchytrium endobioticum</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , & <i>Ditylenchus dipsaci</i> . AD11:- The consignment is free from soil or growth mediums.

- Thalictrum* AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.
- Zantedeschia* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Erwinia chrysanthemi*, & *Phytophthora* spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from *Phymatotrichum omnivorum*, & *Synchytrium endobioticum*. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, & *Ditylenchus dipsaci*. AD11:- The consignment is free from soil or growth mediums.

4.4.3 Whole Plants

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Whole Plants require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

Plants must be **glasshouse grown** and not older than 10 weeks and/or exceeding 200mm in height. All plants grown in the open ground (i.e. not within a glasshouse) require a phytosanitary import permit.

- Abelia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (E. pomi), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Acalypha* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (E. pomi), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Acanthus* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Achillea* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Adenia* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Phytophthora palmivora* did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Adenium* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Phytophthora palmivora* did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Aglaonema* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Erwinia chrysanthemi* & *Verticillium albo-atrum* did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Ajuga* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phyatotrachum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Albertia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phyatotrachum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Alocasia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phyatotrachum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Aloe* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Polyporus sanguineus* did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phyatotrachum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Alstroemeria* AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phyatotrachum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Amaranthaceae* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, Beet curly top virus, Cucumber mosaic virus, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, *Tetranychus viennensis*, Tobacco ringspot virus, Tomato black ring virus, & Tomato ringspot virus. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Amaryllidaceae* AD4:- The consignment is free from *Aceria tulipae*, *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Androsace* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Anigozanthos* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Anthurium* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Radopholus citrophilis*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Aphelandra* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomii*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Aralia* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Verticillium albo-atrum* did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Arbutus* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Ardisia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Arisaema* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Armeria* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Artocarpus* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Koleroga noxia* & *Phytophthora palmivora* did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Asparagus</i>	AD2:- New Zealand is free from <i>Asparagus virus</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Astartea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Azalea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomya rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Baeckea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ballota</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Beaucarnea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza spp.</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Bergenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bignonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Botrychium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bougainvillea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Bouvardia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Brachychiton</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) does not occur in the area of production. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Brighamia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Buddleia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Calathea</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Calceolaria</i>	AD1:- (i) as appears from inspections during the active growth of that <i>Phytophthora</i> spp. did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Callistemon</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Camellia</i>	AD2:- New Zealand is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , and <i>Exobasidium vexans</i> . AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & <i>Xanthomonas gortencovianum</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Campanula</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Cucumber mosaic virus, <i>Puccinia</i> spp., & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & Uredinales. AD6:- The plants comprising the consignment were rooted and grown in sterilised media. and were packed in sterilised media.
<i>Campsis</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cananga</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Capparis</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Carissa</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Caulophyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Ceanothus* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Ceratopetalum* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Chamaelaucium* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Cibotium* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Clerodendron* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Clematis* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Verticillium albo-atrum* did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Cleyera</i>	AD2:- New Zealand is free from <i>Calacarus carinatus</i> , <i>Exobasidium camelliae</i> , and <i>Exobasidium vexans</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Cephaluros parasiticus</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phomopsis theae</i> , <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> & <i>Xanthomonas gortencovianum</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Codonanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Coleus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Columnea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Coprosma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Crossandra</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ctenanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Cussonia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cyathea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Cycadales	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Cyclamen</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Ramularia cyclaminicola</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dahlia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> & fungal wilt diseases did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Damptera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Danae</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dasylyrion</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Delphinium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Beet curly top virus, & Tobacco ringspot virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dendranthema</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma, Chrysanthemum stunt virus, <i>Didymella chrysanthemi</i> , <i>Erwinia chrysanthemi</i> , <i>Puccinia horiana</i> , & Tomato aspermy virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dianthus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, <i>Cacoecimorpha pronubana</i> , Carnation etched ring virus, Carnation necrotic fleck virus, Carnation streak virus, <i>Erwinia chrysanthemi</i> pv. <i>dianthi</i> , Pelargonium leaf curl virus, <i>Pseudomonas caryophylli</i> , ring virus, Carnation necrotic fleck virus, Carnation streak virus, <i>Erwinia chrysanthemi</i> pv. <i>dianthi</i> , Pelargonium leaf curl virus, <i>Pseudomonas caryophylli</i> , & <i>Ustilago violacea</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted and grown in sterilised media. and were packed in sterilised media.

<i>Diascia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dicksonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dieffenbachia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> , & <i>Phytophthora</i> spp., did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> , AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Dipladenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Draba</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Dracaena</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Aspergillus niger f.sp. floridanus</i> & <i>Phyllosticta dracaenae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> , AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Echeveria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Epipremnum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Erica</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora</i> spp., & <i>Verticillium</i> spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Erigeron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Eryngium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Euonymus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Euonymus variegation virus</i> , & <i>Oidium euonymi-japonici</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Euphorbia</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD8:- Culture or tissue culture transplants were obtained from mother plants that were practically free of Euphorbia mosaic virus and Poinsettia mosaic virus.
<i>Eustoma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Exacum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Fatshedera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Fittonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Fuchsia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Gardenia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Genista</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Geogonanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Geranium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Beet curly top virus, Pelargonium leaf curl virus, Tomato black ring virus, <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Gerbera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phytophthora cryptogea</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Grevillea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Gypsophila</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hamamelis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Endothia gyrosa</i> & <i>Polyporus gilvus</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Hebe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hedera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Helichrysum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Helminthostachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Hibbertia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hosta</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hoya</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hydrangea</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Hydrangea ringspot virus, Tobacco ringspot virus & Tomato ringspot virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Hypoestes</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ilex</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Ixora</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Cercospora ixorae</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Jatropha</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phakopsora jatrophiicola</i> & <i>Phytophthora palmivora</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Kalanchoe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Kalmia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lambertia</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media. AD8:- culture or tissue culture transplants were obtained from mother plants that were practically free of Euphorbia mosaic virus and Poinsettia mosaic virus.
<i>Lamium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Lavandula</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Leschenaultia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Limonium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lisianthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lonicera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lychnis</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lycopodium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Lysimachia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Lythrum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Macropidia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Sclerotinia bulborum</i> & <i>Sclerotium tuliparum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , Lily mosaic virus, Lily rosette virus, <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Magnolia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Nectria galligena</i> , <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Marantaceae	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Abaca mosaic virus & <i>Rosselinia bunodes</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Medinilla</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Menianthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Metrosideros</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Mimulus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Peronospora jacksonii</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Mitella</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Monarda</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Monstera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Mussaenda</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Myrmecodia</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nematanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nepenthes</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nertera</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Nolina</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oenanthe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oenothera</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Ophioglossum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Origanum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Osmanthus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Oxalis</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Ennomos subsignarius</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pachypodium</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pachystachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pandanus</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cadang-cadang viroid did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Peltiphyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Penstemon</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Peperomia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Cucumber mosaic virus & Peperomia ringspot virus did not occur on those plants; or those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Petasites</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Puccinia</i> spp., <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Philodendron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Phlox</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pyrenochaeta phlogina</i> & <i>Verticillium albo-atrum</i> did not occur on those plants; or those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomii</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.

<i>Phormium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Phormium</i> yellowing mycoplasma, <i>Uredo phormii</i> & <i>Xanthomonas campestris</i> pv. <i>phormicola</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Physostegia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Phyteuma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pisonia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pittosporum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
Polypodiaceae	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Polyscias</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Preslia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Primula</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Aster yellows mycoplasma did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising this consignment were grown, rooted and packed in sterilised media.
<i>Prunella</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Pseuderanthemum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Psilotum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Pulmonaria* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Punica* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Sphaceloma punicae* did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Pyrenacantha* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Pyrostegia* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Rhadinophora* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Erwinia chrysanthemi* & *Phytophthora* spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Rhaphidiphora* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Erwinia chrysanthemi* & *Phytophthora* spp. did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Rhododendron</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysomyxa rhododendri</i> , <i>Ovulina azaleae</i> & <i>Septoria azaleae</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Rheoe</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ruellia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Pseudomonas solanacearum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Ruscus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Salix</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Physalospora miyabeana</i> , <i>Polaccia saliciperda</i> , <i>Pseudomonas saliciperda</i> , & Virus chlorosis, did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Salvia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

- Sambucus* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned Alfalfa dwarf mosaic virus, Arabis mosaic virus, Cherry leafroll virus & Tobacco ringspot virus did not occur on those plants; or (ii) do not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted and grown in sterilised media. and were packed in sterilised media.
- Sansevieria* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Saxifraga* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Schefflera* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Schismatoglottis* AD1:- (i) as appears from inspections during the active growth of the mother plants concerned *Erwinia chrysanthemi* did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
- Schizaea* AD4:- The consignment is free from *Aphelenchoides besseyi*, *Aphelenchoides fragariae*, *Aphelenchoides ritzema-bosi*, *Ditylenchus destructor*, *Ditylenchus dipsaci*, *Eotetranychus pruni* (*E. pomi*), *Eotetranychus sexmaculatus*, *Frankliniella occidentalis*, *Liriomyza* spp., *Phymatotrichum omnivorum*, *Synchytrium endobioticum*, *Tetranychus canadensis*, *Tetranychus mcdanielli*, *Tetranychus pacificus*, *Tetranychus schoenei*, & *Tetranychus viennensis*. AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Scindapsus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Selaginella</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Simmondsia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Soleirolia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Sonerilla</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Spathiphyllum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stachys</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Statice</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stephanotis</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Chrysanthemum stunt viroid</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> , AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Stereospermum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Strelitzia</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Abaca mosaic virus</i> , <i>Erwinia carotovora</i> pv. <i>musae</i> & <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> did not occur on those plants; or (ii) do not occur in the area of production concerned. AD3:- The area of production is free from <i>Mycosphaerella fijiensis</i> , <i>Pseudomonas solanacearum</i> , & <i>Xanthomonas campestris</i> pv. <i>celebensis</i> . AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Syngonium</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Erwinia chrysanthemi</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> , AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Talinum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Thalictrum</i>	AD4:- The consignment is free from <i>Aceria tulipae</i> , <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tibouchina</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Thymus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tmesipterus</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tolmiea</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Trachelium</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Tradescantia</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

<i>Verbascum</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Veronica</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Viburnum</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Verticillium albo-atrum</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Wisteria</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Xanthosoma</i>	AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.
<i>Yucca</i>	AD1:- (i) as appears from inspections during the active growth of the mother plants concerned <i>Leptosphaeria obtusispora</i> did not occur on those plants; or (ii) does not occur in the area of production concerned. AD4:- The consignment is free from <i>Aphelenchoides besseyi</i> , <i>Aphelenchoides fragariae</i> , <i>Aphelenchoides ritzema-bosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Eotetranychus pruni</i> (<i>E. pomi</i>), <i>Eotetranychus sexmaculatus</i> , <i>Frankliniella occidentalis</i> , <i>Liriomyza</i> spp., <i>Phymatotrichum omnivorum</i> , <i>Synchytrium endobioticum</i> , <i>Tetranychus canadensis</i> , <i>Tetranychus mcdanielli</i> , <i>Tetranychus pacificus</i> , <i>Tetranychus schoenei</i> , & <i>Tetranychus viennensis</i> . AD6:- The plants comprising the consignment were rooted, grown and packed in sterilised media.

4.4.4 Tissue Culture *in vitro*

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Tissue Culture *in vitro* require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Achillea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Actinidia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aglaonema</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Alocasia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Amaranthaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Cucumber mosaic virus, Tobacco ringspot virus, Tomato black ring virus & Tomato ringspot virus.
Amaryllidaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Anigozanthos</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Anthurium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aphelandra</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Aralia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ardisia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Asparagus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Asparagus virus.
<i>Aster</i>	AD4:- The consignment is free Uredinales (Pine rusts). AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, Tobacco rattle virus, & Tomato aspermy virus.
<i>Beaucarnea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Begonia</i> (excluding bulbous varieties)	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Botrychium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Bouvardia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Brachychiton</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Bromeliaceae (excluding <i>Ananas</i> spp.)	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Calathea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Campanula</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Cucumber mosaic virus,
<i>Ceropegia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Chamaelaucium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Clematis</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Clerodendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Codonanthe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Columnea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cordyline</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Crossandra</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ctenanthe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cupressus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Cussonia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dahlia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Dendranthema</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, Chrysanthemum stunt viroid, & Tomato aspermy virus.
<i>Dianthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Carnation etched ring virus, Carnation necrotic fleck virus, Carnation streak virus, & Pelargonium leaf curl virus.
<i>Dieffenbachia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dipladenia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Draba</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Dracaena</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Echeveria</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Erica</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Euphorbia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Euphorbia mosaic virus & Poinsettia mosaic virus.
<i>Fuchsia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Alfalfa dwarf mosaic virus.
<i>Geogenanthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Geranium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, Pelargonium leaf curl virus, & Tomato black ring virus.
<i>Gerbera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Grevillea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Gypsophila</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hedera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Alfalfa dwarf mosaic virus.

<i>Heliconia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus & Banana bunchy top virus.
<i>Helminthostachys</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hibiscus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Beet curly top virus, & Cotton leaf curl virus.
<i>Hoya</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Hydrangea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Hydrangea ringspot virus, Tobacco ringspot virus, & Tomato ringspot virus.
<i>Kalanchoe</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Kalmia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lambertia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Euphorbia mosaic virus & Poinsettia mosaic virus.
<i>Limonium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lisianthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lycopodium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Lysimachia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Marantaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus, & <i>Rosselinia bunodes</i> .
<i>Monstera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Nematanthus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Nertera</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ophioglossum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Ophiopogon</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Orchidaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Pachypodium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Passiflora</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Penstemon</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Cucumber mosaic virus.
<i>Philodendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Phlox</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Tomato aspermy virus.
<i>Phyloglossum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Phyteuma</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Pisonia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
Polypodiaceae	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Polyscias</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Psilotum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Rhododendron</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Ruscus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Sansevieria</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Saxifraga</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Schizaea</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Scindapsus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Simmondsia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

<i>Solidaster</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Aster yellows mycoplasma, & Tobacco rattle virus.
<i>Sonerilla</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Spathiphyllum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Stephanotis</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Chrysanthemum stunt viroid.
<i>Strelitzia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases. AD10:- The mother plants from which the tissue or tissue culture comprising the consignment were obtained, were indexed for and found free from Abaca mosaic virus & Banana bunchy top virus.
<i>Syngonium</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Thalictrum</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tibouchina</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tmesipterus</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Tradescantia</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.
<i>Yucca</i>	AD8:- The tissue culture was obtained from mother plants that are practically free of viruses and other diseases.

4.5 Seeds (Grain)/Nuts

4.5.1 Seed for Sowing

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Seed for Sowing require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Abrus</i>	No additional declarations required.
<i>Abutilon</i>	No additional declarations required.
<i>Acaena</i>	Phytosanitary certificate not required.
<i>Acanthus</i>	Phytosanitary certificate not required.
<i>Achillea</i>	Phytosanitary certificate not required.
<i>Achimenes</i>	Phytosanitary certificate not required.
<i>Aciphylla</i>	Phytosanitary certificate not required.
<i>Aconitum</i>	Phytosanitary certificate not required.
<i>Acokanthera</i>	No additional declarations required.
<i>Acrocarpus</i>	No additional declarations required.
<i>Actaea</i>	Phytosanitary certificate not required.
<i>Actinidia</i>	Phytosanitary certificate not required.
<i>Actinotus</i>	Phytosanitary certificate not required.
<i>Adansonia</i>	No additional declarations required.
<i>Adenophora</i>	Phytosanitary certificate not required.
<i>Adenostyles</i>	Phytosanitary certificate not required.
<i>Adenanthos</i>	No additional declarations required.
<i>Adenium</i>	No additional declarations required.
<i>Aethionema</i>	Phytosanitary certificate not required.
<i>Aetoxicon</i>	No additional declarations required.
<i>Agastache</i>	Phytosanitary certificate not required.
<i>Agathis</i>	No additional declarations required.
<i>Agave</i>	No additional declarations required.
<i>Ageratum</i>	Phytosanitary certificate not required.
<i>Aglaonema</i>	Phytosanitary certificate not required.

<i>Agrostemma</i>	Phytosanitary certificate not required.
<i>Agonis</i>	No additional declarations required.
<i>Akebia</i>	No additional declarations required.
<i>Alberta</i>	No additional declarations required.
<i>Alcea</i>	Phytosanitary certificate not required.
<i>Alectryon</i>	No additional declarations required.
<i>Aleurites</i>	No additional declarations required.
<i>Alliaria</i>	Phytosanitary certificate not required.
<i>Alloteropsis</i>	No additional declarations required.
<i>Alluaudia</i>	No additional declarations required.
<i>Alnus</i>	No additional declarations required.
<i>Alocasia</i>	No additional declarations required.
<i>Aloe</i>	No additional declarations required.
<i>Aloysia</i>	No additional declarations required.
<i>Alseuosmia</i>	Phytosanitary certificate not required.
<i>Alstroemeria</i>	Phytosanitary certificate not required.
<i>Althaea</i>	Phytosanitary certificate not required.
<i>Alyssoides</i>	Phytosanitary certificate not required.
<i>Alyssum</i>	Phytosanitary certificate not required.
<i>Amaranthus</i>	Phytosanitary certificate not required.
<i>Amaryllis</i>	Phytosanitary certificate not required.
<i>Ammi</i>	Phytosanitary certificate not required.
<i>Ammobium</i>	Phytosanitary certificate not required.
<i>Amorpha</i>	No additional declarations required.
<i>Amorphophallus</i>	Phytosanitary certificate not required.
<i>Amphicarpea</i>	No additional declarations required.
<i>Amsonia</i>	Phytosanitary certificate not required.
<i>Anacyclus</i>	Phytosanitary certificate not required.
<i>Ananas</i>	Phytosanitary certificate not required.
<i>Anaphalis</i>	Phytosanitary certificate not required.
<i>Anchusa</i>	Phytosanitary certificate not required.
<i>Angophora</i>	No additional declarations required.

<i>Anemone</i>	Phytosanitary certificate not required.
<i>Anethum</i>	Phytosanitary certificate not required.
<i>Angelica</i>	Phytosanitary certificate not required.
<i>Anigozanthos</i>	Phytosanitary certificate not required.
<i>Anthocleista</i>	No additional declarations required.
<i>Antennaria</i>	Phytosanitary certificate not required.
<i>Anthemis</i>	Phytosanitary certificate not required.
<i>Antholyza</i>	Phytosanitary certificate not required.
<i>Anthriscus</i>	Phytosanitary certificate not required.
<i>Anthurium</i>	Phytosanitary certificate not required.
<i>Anthyllis</i>	Phytosanitary certificate not required.
<i>Antidesma</i>	No additional declarations required.
<i>Antigonon</i>	No additional declarations required.
<i>Antirrhinum</i>	Phytosanitary certificate not required.
<i>Aphelandra</i>	Phytosanitary certificate not required.
<i>Apios</i>	Phytosanitary certificate not required.
<i>Apocynum</i>	Phytosanitary certificate not required.
<i>Aquilegia</i>	Phytosanitary certificate not required.
<i>Aralia</i>	No additional declarations required.
<i>Arbutus</i>	No additional declarations required.
<i>Arctium</i>	Phytosanitary certificate not required.
<i>Arctotis</i>	Phytosanitary certificate not required.
<i>Arctostaphylos</i>	No additional declarations required.
<i>Ardisa</i>	No additional declarations required.
<i>Argyroderma</i>	Phytosanitary certificate not required.
<i>Arisaema</i>	No additional declarations required.
<i>Aristolochia</i>	No additional declarations required.
<i>Armeria</i>	Phytosanitary certificate not required.
<i>Artemisia</i>	Phytosanitary certificate not required.
<i>Artocarpus</i>	No additional declarations required.
<i>Arum</i>	No additional declarations required.
<i>Asarina</i>	Phytosanitary certificate not required.

<i>Asclepias</i>	Phytosanitary certificate not required.
<i>Asimina</i>	No additional declarations required.
<i>Astartea</i>	No additional declarations required.
<i>Aster</i>	Phytosanitary certificate not required.
<i>Astragalus</i>	Phytosanitary certificate not required.
<i>Astrantia</i>	Phytosanitary certificate not required.
<i>Asyneuma</i>	Phytosanitary certificate not required.
<i>Atalaya</i>	No additional declarations required.
<i>Atriplex</i>	No additional declarations required.
<i>Aubrieta</i>	Phytosanitary certificate not required.
<i>Aurinia</i>	Phytosanitary certificate not required.
<i>Axonopus</i>	Phytosanitary certificate not required.
<i>Azadirachta</i>	No additional declarations required.
<i>Azanza</i>	No additional declarations required.
<i>Azara</i>	No additional declarations required.
<i>Baeckea</i>	No additional declarations required.
<i>Balanites</i>	No additional declarations required.
<i>Ballota</i>	Phytosanitary certificate not required.
<i>Balsamita</i>	Phytosanitary certificate not required.
<i>Banksia</i>	No additional declarations required.
<i>Baptisia</i>	Phytosanitary certificate not required.
<i>Basilicum</i>	Phytosanitary certificate not required.
<i>Bassia</i>	Phytosanitary certificate not required.
<i>Bauhinia</i>	No additional declarations required.
<i>Beaucarnea</i>	No additional declarations required.
<i>Beaufortia</i>	No additional declarations required.
<i>Beaumontia</i>	No additional declarations required.
<i>Begonia</i>	Phytosanitary certificate not required.
<i>Bergenia</i>	Phytosanitary certificate not required.
<i>Berlandiera</i>	Phytosanitary certificate not required.
<i>Beschorneria</i>	Phytosanitary certificate not required.
<i>Bignonia</i>	No additional declarations required.

<i>Bischofia</i>	No additional declarations required.
<i>Biserrula</i>	No additional declarations required.
<i>Boltonia</i>	Phytosanitary certificate not required.
<i>Borago</i>	Phytosanitary certificate not required.
<i>Bossiaea</i>	No additional declarations required.
<i>Botrychium</i>	No additional declarations required.
<i>Bougainvillea</i>	No additional declarations required.
<i>Bouvardia</i>	Phytosanitary certificate not required.
<i>Brachychiton</i>	No additional declarations required.
<i>Brachycome</i>	Phytosanitary certificate not required.
<i>Brachylaena</i>	No additional declarations required.
<i>Brachysema</i>	Phytosanitary certificate not required.
<i>Brachystelma</i>	Phytosanitary certificate not required.
<i>Brachystemma</i>	No additional declarations required.
<i>Brachystegia</i>	Phytosanitary certificate not required.
<i>Bridelia</i>	No additional declarations required.
<i>Brighamia</i>	No additional declarations required.
<i>Brizia</i>	No additional declarations required.
<i>Bromelia</i>	Phytosanitary certificate not required.
<i>Brownea</i>	No additional declarations required.
<i>Browallia</i>	Phytosanitary certificate not required.
<i>Brugmansia</i>	Phytosanitary certificate not required.
<i>Brunfelsia</i>	No additional declarations required.
<i>Buckinghamia</i>	No additional declarations required.
<i>Buddleia</i>	No additional declarations required.
<i>Bupleurum</i>	No additional declarations required.
<i>Buphthalmum</i>	Phytosanitary certificate not required.
<i>Burchellia</i>	No additional declarations required.
<i>Bursaria</i>	No additional declarations required.
<i>Bursera</i>	No additional declarations required.
<i>Byblis</i>	Phytosanitary certificate not required.
<i>Cajanus</i>	No additional declarations required.

<i>Caladium</i>	No additional declarations required.
<i>Calamintha</i>	Phytosanitary certificate not required.
<i>Calamus</i>	No additional declarations required.
<i>Calathea</i>	Phytosanitary certificate not required.
<i>Calceolaria</i>	Phytosanitary certificate not required.
<i>Calendula</i>	Phytosanitary certificate not required.
<i>Calliandra</i>	No additional declarations required.
<i>Callistemon</i>	No additional declarations required.
<i>Callistephus</i>	Phytosanitary certificate not required.
<i>Callitris</i>	No additional declarations required.
<i>Calocephalus</i>	Phytosanitary certificate not required.
<i>Calothamnus</i>	No additional declarations required.
<i>Calpurnia</i>	No additional declarations required.
<i>Caltha</i>	Phytosanitary certificate not required.
<i>Calycanthus</i>	Phytosanitary certificate not required.
<i>Campanula</i>	Phytosanitary certificate not required.
<i>Cananga</i>	No additional declarations required.
<i>Canavalia</i>	Phytosanitary certificate not required.
<i>Canna</i>	Phytosanitary certificate not required.
<i>Canthium</i>	No additional declarations required.
<i>Capparis</i>	No additional declarations required.
<i>Carangana</i>	No additional declarations required.
<i>Caralluma</i>	Phytosanitary certificate not required.
<i>Carissa</i>	No additional declarations required.
<i>Carlina</i>	Phytosanitary certificate not required.
<i>Carludovica</i>	Phytosanitary certificate not required.
<i>Carpanthea</i>	Phytosanitary certificate not required.
<i>Carpinus</i>	No additional declarations required.
<i>Carpodetus</i>	No additional declarations required.
<i>Carthamus</i>	Phytosanitary certificate not required.
<i>Caryocar</i>	No additional declarations required.
<i>Caryopteris</i>	No additional declarations required.

<i>Cassia</i>	No additional declarations required.
<i>Cassine</i>	Phytosanitary certificate not required.
<i>Cassiope</i>	No additional declarations required.
<i>Catanache</i>	Phytosanitary certificate not required.
<i>Catharanthus</i>	Phytosanitary certificate not required.
<i>Ceanothus</i>	No additional declarations required.
<i>Cecropia</i>	No additional declarations required.
<i>Cedrela</i>	No additional declarations required.
<i>Cedrus</i>	No additional declarations required.
<i>Celosia</i>	Phytosanitary certificate not required.
<i>Celtis</i>	No additional declarations required.
<i>Cenchrus</i>	No additional declarations required.
<i>Centaurea</i>	Phytosanitary certificate not required.
<i>Centranthus</i>	Phytosanitary certificate not required.
<i>Cephalaria</i>	Phytosanitary certificate not required.
<i>Cephalotaxus</i>	No additional declarations required.
<i>Cerastium</i>	Phytosanitary certificate not required.
<i>Ceratonia</i>	No additional declarations required.
<i>Ceratopetalum</i>	No additional declarations required.
<i>Cercidium</i>	No additional declarations required.
<i>Cerintho</i>	Phytosanitary certificate not required.
<i>Ceropegia</i>	Phytosanitary certificate not required.
<i>Cercis</i>	No additional declarations required.
<i>Chamaecytisus</i>	No additional declarations required.
<i>Chamaelaucium</i>	No additional declarations required.
<i>Cheiranthus</i>	Phytosanitary certificate not required.
<i>Chelone</i>	Phytosanitary certificate not required.
<i>Chionanthus</i>	No additional declarations required.
<i>Chionochoa</i>	No additional declarations required.
<i>Chloris</i>	No additional declarations required.
<i>Chorizema</i>	No additional declarations required.
<i>Chrysophyllum</i>	No additional declarations required.

<i>Chrysopsis</i>	Phytosanitary certificate not required.
<i>Chukrasia</i>	No additional declarations required.
<i>Cichorium</i>	No additional declarations required.
<i>Cimicifuga</i>	Phytosanitary certificate not required.
<i>Cineraria</i>	Phytosanitary certificate not required.
<i>Cinnamomum</i>	No additional declarations required.
<i>Cistus</i>	No additional declarations required.
<i>Cladanthus</i>	Phytosanitary certificate not required.
<i>Cladrastis</i>	No additional declarations required.
<i>Clarkia</i>	Phytosanitary certificate not required.
<i>Cleome</i>	Phytosanitary certificate not required.
<i>Clerodendrum</i>	No additional declarations required.
<i>Clianthus</i>	No additional declarations required.
<i>Clinopodium</i>	Phytosanitary certificate not required.
<i>Clitoria</i>	No additional declarations required.
<i>Clivia</i>	Phytosanitary certificate not required.
<i>Cneorum</i>	No additional declarations required.
<i>Cobaea</i>	No additional declarations required.
<i>Coccoloba</i>	No additional declarations required.
<i>Cochlearia</i>	Phytosanitary certificate not required.
<i>Cochlospermum</i>	No additional declarations required.
<i>Codiaeum</i>	No additional declarations required.
<i>Codonopsis</i>	Phytosanitary certificate not required.
<i>Cola</i>	No additional declarations required.
<i>Colchicum</i>	No additional declarations required.
<i>Coleus</i>	Phytosanitary certificate not required.
<i>Collinsia</i>	Phytosanitary certificate not required.
<i>Colubrina</i>	No additional declarations required.
<i>Colutea</i>	No additional declarations required.
<i>Colvillea</i>	No additional declarations required.
<i>Commiphora</i>	Phytosanitary certificate not required.
<i>Conophytum</i>	Phytosanitary certificate not required.

<i>Consolida</i>	Phytosanitary certificate not required.
<i>Convolvulus</i>	Phytosanitary certificate not required.
<i>Coprosma</i>	No additional declarations required.
<i>Cordyline</i>	No additional declarations required.
<i>Coriandrum</i>	Phytosanitary certificate not required.
<i>Corema</i>	No additional declarations required.
<i>Coriaria</i>	No additional declarations required.
<i>Coreopsis</i>	Phytosanitary certificate not required.
<i>Coronilla</i>	No additional declarations required.
<i>Cortaderia</i>	No additional declarations required.
<i>Corydalis</i>	No additional declarations required.
<i>Corynocarpus</i>	No additional declarations required.
<i>Cosmos</i>	Phytosanitary certificate not required.
<i>Cotinus</i>	No additional declarations required.
<i>Cotoneaster</i>	No additional declarations required.
<i>Crambe</i>	Phytosanitary certificate not required.
<i>Crepis</i>	Phytosanitary certificate not required.
<i>Crescentia</i>	Phytosanitary certificate not required.
<i>Crinodendron</i>	No additional declarations required.
<i>Crossandra</i>	Phytosanitary certificate not required.
<i>Croton</i>	No additional declarations required.
<i>Cryptotaenia</i>	Phytosanitary certificate not required.
<i>Cuminum</i>	Phytosanitary certificate not required.
<i>Cupaniopsis</i>	No additional declarations required.
<i>Cuphea</i>	Phytosanitary certificate not required.
<i>Curculigo</i>	Phytosanitary certificate not required.
<i>Curcuma</i>	Phytosanitary certificate not required.
<i>Cussonia</i>	No additional declarations required.
<i>Cyathea</i>	No additional declarations required.
Cycadales	No additional declarations required.
<i>Cyclamen</i>	Phytosanitary certificate not required.
<i>Cymbopogon</i>	No additional declarations required.

<i>Cynara</i>	Phytosanitary certificate not required.
<i>Cynodon</i>	No additional declarations required.
<i>Cynoglossum</i>	Phytosanitary certificate not required.
<i>Cytisus</i>	No additional declarations required.
<i>Dahlia</i>	Phytosanitary certificate not required.
<i>Dais</i>	No additional declarations required.
<i>Dalbergia</i>	No additional declarations required.
<i>Danae</i>	No additional declarations required.
<i>Darlingtonia</i>	Phytosanitary certificate not required.
<i>Daphne</i>	No additional declarations required.
<i>Darwinia</i>	No additional declarations required.
<i>Dasyllirion</i>	No additional declarations required.
<i>Davidia</i>	No additional declarations required.
<i>Delonix</i>	No additional declarations required.
<i>Diascia</i>	Phytosanitary certificate not required.
<i>Dialium</i>	No additional declarations required.
<i>Dicentra</i>	Phytosanitary certificate not required.
<i>Dichondra</i>	Phytosanitary certificate not required.
<i>Didelta</i>	Phytosanitary certificate not required.
<i>Didierea</i>	No additional declarations required.
<i>Didymaotus</i>	No additional declarations required.
<i>Dieffenbachia</i>	No additional declarations required.
<i>Dillenia</i>	No additional declarations required.
<i>Dimocarpus</i>	No additional declarations required.
<i>Dimorphotheca</i>	Phytosanitary certificate not required.
<i>Dionaea</i>	Phytosanitary certificate not required.
<i>Dioscorea</i>	Phytosanitary certificate not required.
<i>Dipsacus</i>	Phytosanitary certificate not required.
<i>Dipterocarpus</i>	No additional declarations required.
<i>Dischidia</i>	Phytosanitary certificate not required.
<i>Dodecatheon</i>	Phytosanitary certificate not required.
<i>Dodonaea</i>	No additional declarations required.

<i>Dombeya</i>	No additional declarations required.
<i>Doronicum</i>	Phytosanitary certificate not required.
<i>Dorstenia</i>	No additional declarations required.
<i>Doryanthes</i>	Phytosanitary certificate not required.
<i>Draba</i>	Phytosanitary certificate not required.
<i>Dracaena</i>	No additional declarations required.
<i>Dracocephalum</i>	Phytosanitary certificate not required.
<i>Drosera</i>	Phytosanitary certificate not required.
<i>Dryandra</i>	No additional declarations required.
<i>Dysoxylum</i>	No additional declarations required.
<i>Duvalia</i>	Phytosanitary certificate not required.
<i>Echeveria</i>	Phytosanitary certificate not required.
<i>Echinacea</i>	Phytosanitary certificate not required.
<i>Echinops</i>	Phytosanitary certificate not required.
<i>Echium</i>	Phytosanitary certificate not required.
<i>Ehretia</i>	No additional declarations required.
<i>Elacocarpus</i>	No additional declarations required.
<i>Elaeagnus</i>	No additional declarations required.
<i>Elettaria</i>	Phytosanitary certificate not required.
<i>Eleutherococcus</i>	No additional declarations required.
<i>Edgeworthia</i>	No additional declarations required.
<i>Enterolobium</i>	No additional declarations required.
<i>Epacris</i>	No additional declarations required.
<i>EphedraEpilobium</i>	No additional declarations required. Phytosanitary certificate not required.
<i>Eremophila</i>	No additional declarations required.
<i>Erigeron</i>	Phytosanitary certificate not required.
<i>Eriobotrya</i>	No additional declarations required.
<i>Eriogonum</i>	Phytosanitary certificate not required.
<i>Eriophyllum</i>	No additional declarations required.
<i>Eruca</i>	Phytosanitary certificate not required.
<i>Eryngium</i>	Phytosanitary certificate not required.
<i>Erysimum</i>	Phytosanitary certificate not required.

<i>Erythrina</i>	No additional declarations required.
<i>Escallonia</i>	No additional declarations required.
<i>Eschscholzia</i>	Phytosanitary certificate not required.
<i>Euclea</i>	No additional declarations required.
<i>Eucommia</i>	No additional declarations required.
<i>Eucryphia</i>	No additional declarations required.
<i>Eupatorium</i>	Phytosanitary certificate not required.
<i>Euphorbia</i>	No additional declarations required.
<i>Euryops</i>	Phytosanitary certificate not required.
<i>Eustoma</i>	Phytosanitary certificate not required.
<i>Euptelea</i>	No additional declarations required.
<i>Evolvulus</i>	Phytosanitary certificate not required.
<i>Exacum</i>	Phytosanitary certificate not required.
<i>Faucaria</i>	No additional declarations required.
<i>Felicia</i>	Phytosanitary certificate not required.
<i>Ficus</i>	No additional declarations required.
<i>Fittonia</i>	Phytosanitary certificate not required.
<i>Flacourtia</i>	No additional declarations required.
<i>Flindersia</i>	No additional declarations required.
<i>Foeniculum</i>	Phytosanitary certificate not required.
<i>Fremontodenron</i>	No additional declarations required.
<i>Freesia</i>	Phytosanitary certificate not required.
<i>Fuchsia</i>	Phytosanitary certificate not required.
<i>Fumaria</i>	Phytosanitary certificate not required.
<i>Furcraea</i>	No additional declarations required.
<i>Gahnia</i>	No additional declarations required.
<i>Gaillardia</i>	No additional declarations required.
<i>Galega</i>	Phytosanitary certificate not required.
<i>Galium</i>	Phytosanitary certificate not required.
<i>Galphimia</i>	No additional declarations required.
<i>Gardenia</i>	No additional declarations required.
<i>Garcinia</i>	No additional declarations required.

<i>Gaura</i>	Phytosanitary certificate not required.
<i>Gazania</i>	Phytosanitary certificate not required.
<i>Genista</i>	No additional declarations required.
<i>Gentiana</i>	Phytosanitary certificate not required.
<i>Geranium</i>	Phytosanitary certificate not required.
<i>Gerbera</i>	Phytosanitary certificate not required.
<i>Geum</i>	Phytosanitary certificate not required.
<i>Gilia</i>	Phytosanitary certificate not required.
<i>Ginkgo</i>	No additional declarations required.
<i>Gladiolus</i>	Phytosanitary certificate not required.
<i>Glaucium</i>	Phytosanitary certificate not required.
<i>Gleditsia</i>	No additional declarations required.
<i>Gliricidia</i>	No additional declarations required.
<i>Globularia</i>	Phytosanitary certificate not required.
<i>Gloxinia</i>	Phytosanitary certificate not required.
<i>Gmelina</i>	No additional declarations required.
<i>Godetia</i>	Phytosanitary certificate not required.
<i>Gomphrena</i>	Phytosanitary certificate not required.
<i>Goodenia</i>	No additional declarations required.
<i>Gossypium</i>	No additional declarations required. Must be acid de-linted.
<i>Gosswilerodendron</i>	No additional declarations required.
<i>Gouania</i>	No additional declarations required.
<i>Grevillea</i>	No additional declarations required.
<i>Greyia</i>	No additional declarations required.
<i>Grindelia</i>	Phytosanitary certificate not required.
<i>Griselina</i>	No additional declarations required.
<i>Gymnosporia</i>	No additional declarations required.
<i>Gypsophila</i>	Phytosanitary certificate not required.
<i>Haloxylon</i>	No additional declarations required.
<i>Hamamelis</i>	No additional declarations required.
<i>Hardenbergia</i>	No additional declarations required.
<i>Harungana</i>	No additional declarations required.

<i>Hebe</i>	No additional declarations required.
<i>Hedera</i>	Phytosanitary certificate not required.
<i>Hebenstretia</i>	No additional declarations required.
<i>Hedysarum</i>	No additional declarations required.
<i>Helenium</i>	Phytosanitary certificate not required.
<i>Helichrysum</i>	Phytosanitary certificate not required.
<i>Helinus</i>	No additional declarations required.
<i>Heliophila</i>	Phytosanitary certificate not required.
<i>Heliopsis</i>	Phytosanitary certificate not required.
<i>Heliocarpus</i>	No additional declarations required.
<i>Helleborus</i>	No additional declarations required.
<i>Helminthostachys</i>	No additional declarations required.
<i>Hemigenia</i>	No additional declarations required.
<i>Heliotropium</i>	Phytosanitary certificate not required.
<i>Helipterum</i>	Phytosanitary certificate not required.
<i>Hesperis</i>	Phytosanitary certificate not required.
<i>Heuchera</i>	Phytosanitary certificate not required.
<i>Hibbertia</i>	No additional declarations required.
<i>Hieracium</i>	Phytosanitary certificate not required.
<i>Hildegardia</i>	No additional declarations required.
<i>Hippophae</i>	No additional declarations required.
<i>Hoheria</i>	No additional declarations required.
<i>Hosta</i>	Phytosanitary certificate not required.
<i>Hovea</i>	No additional declarations required.
<i>Hovenia</i>	No additional declarations required.
<i>Hyacinthus</i>	Phytosanitary certificate not required.
<i>Hydrocotyle</i>	Phytosanitary certificate not required.
<i>Hymenachne</i>	No additional declarations required.
<i>Hymenanthrum</i>	No additional declarations required.
<i>Hymenosporum</i>	No additional declarations required.
<i>Hyoscyamus</i>	Phytosanitary certificate not required.
<i>Hypocalymma</i>	No additional declarations required.

<i>Hypoestes</i>	Phytosanitary certificate not required.
<i>Hyssopus</i>	Phytosanitary certificate not required.
<i>Iberis</i>	Phytosanitary certificate not required.
<i>Ilex</i>	No additional declarations required.
<i>Impatiens</i>	No additional declarations required.
<i>Incarvillea</i>	Phytosanitary certificate not required.
<i>Indigofera</i>	No additional declarations required.
<i>Inula</i>	Phytosanitary certificate not required.
<i>Isatis</i>	Phytosanitary certificate not required.
<i>Ixodia</i>	Phytosanitary certificate not required.
<i>Ixora</i>	No additional declarations required.
<i>Jasione</i>	Phytosanitary certificate not required.
<i>Jatropha</i>	No additional declarations required.
<i>Justica</i>	No additional declarations required.
<i>Jurinea</i>	Phytosanitary certificate not required.
<i>Kalanchoe</i>	Phytosanitary certificate not required.
<i>Kalmia</i>	No additional declarations required.
<i>Kennedia</i>	No additional declarations required.
<i>Khaya</i>	No additional declarations required.
<i>Kiggelaria</i>	No additional declarations required.
<i>Kirengeshoma</i>	No additional declarations required.
<i>Kirkia</i>	No additional declarations required.
<i>Kissenia</i>	No additional declarations required.
<i>Knautia</i>	Phytosanitary certificate not required.
<i>Knightsia</i>	No additional declarations required.
<i>Kniphofia</i>	Phytosanitary certificate not required.
<i>Kochia</i>	Phytosanitary certificate not required.
<i>Koelreuteria</i>	No additional declarations required.
<i>Kolkritzia</i>	No additional declarations required.
<i>Kunzea</i>	No additional declarations required.
<i>Laburnum</i>	No additional declarations required.
<i>Lachnostachys</i>	Phytosanitary certificate not required.

<i>Lagerstroemia</i>	No additional declarations required.
<i>Landolphia</i>	No additional declarations required.
<i>Langsdorffia</i>	No additional declarations required.
<i>Lannea</i>	No additional declarations required.
<i>Laportea</i>	No additional declarations required.
<i>Lasthenia</i>	Phytosanitary certificate not required.
<i>Lathyrus</i>	Phytosanitary certificate not required.
<i>Larix</i>	No additional declarations required.
<i>Laurelia</i>	No additional declarations required.
<i>Laurentia</i>	Phytosanitary certificate not required.
<i>Laurus</i>	No additional declarations required.
<i>Lavandula</i>	Phytosanitary certificate not required.
<i>Lavatera</i>	Phytosanitary certificate not required.
<i>Layia</i>	Phytosanitary certificate not required.
<i>Leonotis</i>	Phytosanitary certificate not required.
<i>Leontopodium</i>	Phytosanitary certificate not required.
<i>Lepidium</i>	Phytosanitary certificate not required.
<i>Lechenaultia</i>	No additional declarations required.
<i>Lespedeza</i>	Phytosanitary certificate not required.
<i>Lesquerella</i>	No additional declarations required.
<i>Leucanthemum</i>	Phytosanitary certificate not required.
<i>Leucopogon</i>	No additional declarations required.
<i>Leucothoe</i>	No additional declarations required.
<i>Levisticum</i>	Phytosanitary certificate not required.
<i>Lewisia</i>	Phytosanitary certificate not required.
<i>Liatris</i>	Phytosanitary certificate not required.
<i>Ligularia</i>	Phytosanitary certificate not required.
<i>Lilium</i>	Phytosanitary certificate not required.
<i>Limnanthes</i>	Phytosanitary certificate not required.
<i>Limonium</i>	Phytosanitary certificate not required.
<i>Linaria</i>	Phytosanitary certificate not required.
<i>Lindera</i>	No additional declarations required.

<i>Lisianthus</i>	Phytosanitary certificate not required.
<i>Liquidambar</i>	No additional declarations required.
<i>Litchi</i>	No additional declarations required.
<i>Lithops</i>	No additional declarations required.
<i>Lobelia</i>	No additional declarations required.
<i>Lobularia</i>	Phytosanitary certificate not required.
<i>Lomatium</i>	Phytosanitary certificate not required.
<i>Lonas</i>	Phytosanitary certificate not required.
<i>Lophiocarpus</i>	No additional declarations required.
<i>Lophotocarpus</i>	No additional declarations required.
<i>Lotus</i>	No additional declarations required.
<i>Luculia</i>	No additional declarations required.
<i>Luehea</i>	No additional declarations required.
<i>Lunaria</i>	Phytosanitary certificate not required.
<i>Luzula</i>	Phytosanitary certificate not required.
<i>Lychnis</i>	Phytosanitary certificate not required.
<i>Lycopodiella</i>	Phytosanitary certificate not required.
<i>Lycopodium</i>	Phytosanitary certificate not required.
<i>Lycopus</i>	Phytosanitary certificate not required.
<i>Lysichiton</i>	Phytosanitary certificate not required.
<i>Lysimachia</i>	Phytosanitary certificate not required.
<i>Lythrum</i>	Phytosanitary certificate not required.
<i>Macfadyena</i>	No additional declarations required.
<i>Macarthuria</i>	Phytosanitary certificate not required.
<i>Macleaya</i>	Phytosanitary certificate not required.
<i>Macropidia</i>	Phytosanitary certificate not required.
<i>Macropiper</i>	No additional declarations required.
<i>Macroptilium</i>	No additional declarations required.
<i>Maesa</i>	No additional declarations required.
<i>Magnolia</i>	No additional declarations required.
<i>Malcolmia</i>	Phytosanitary certificate not required.
<i>Malva</i>	Phytosanitary certificate not required.

<i>Mammea</i>	No additional declarations required.
<i>Mandevilla</i>	No additional declarations required.
<i>Manilkara</i>	No additional declarations required.
<i>Maranta</i>	Phytosanitary certificate not required.
<i>Markhamia</i>	No additional declarations required.
<i>Marrubium</i>	No additional declarations required.
<i>Matricaria</i>	Phytosanitary certificate not required.
<i>Matthiola</i>	Phytosanitary certificate not required.
<i>Medinilla</i>	No additional declarations required.
<i>Meconopsis</i>	Phytosanitary certificate not required.
<i>Melaleuca</i>	No additional declarations required.
<i>Medusagyne</i>	Phytosanitary certificate not required.
<i>Melampodium</i>	Phytosanitary certificate not required.
<i>Melianthus</i>	No additional declarations required.
<i>Melissa</i>	Phytosanitary certificate not required.
<i>Memecylon</i>	No additional declarations required.
<i>Mentzelia</i>	No additional declarations required.
<i>Menyanthes</i>	Phytosanitary certificate not required.
<i>Mertensia</i>	Phytosanitary certificate not required.
<i>Meryta</i>	No additional declarations required.
<i>Mesembryanthemum</i>	Phytosanitary certificate not required.
<i>Metrosideros</i>	No additional declarations required.
<i>Meum</i>	No additional declarations required.
<i>Miconia</i>	No additional declarations required.
<i>Mikania</i>	No additional declarations required.
<i>Millettia</i>	No additional declarations required.
<i>Mimosa</i>	No additional declarations required.
<i>Mimulus</i>	Phytosanitary certificate not required.
<i>Mimusops</i>	No additional declarations required.
<i>Mina</i>	Phytosanitary certificate not required.
<i>Momordica</i>	No additional declarations required.

<i>Mirabilis</i>	Phytosanitary certificate not required.
<i>Moluccella</i>	Phytosanitary certificate not required.
<i>Monarda</i>	Phytosanitary certificate not required.
<i>Monstera</i>	Phytosanitary certificate not required.
<i>Morina</i>	Phytosanitary certificate not required.
<i>Moringa</i>	No additional declarations required.
<i>Mundulea</i>	Phytosanitary certificate not required.
<i>Mucuna</i>	No additional declarations required.
<i>Myosotidium</i>	Phytosanitary certificate not required.
<i>Myristica</i>	No additional declarations required.
<i>Myrrhis</i>	Phytosanitary certificate not required.
<i>Myrsine</i>	No additional declarations required.
<i>Myrtus</i>	No additional declarations required.
<i>Nandina</i>	Phytosanitary certificate not required.
<i>Nasturtium</i>	Phytosanitary certificate not required.
<i>Nemesia</i>	Phytosanitary certificate not required.
<i>Nemophila</i>	Phytosanitary certificate not required.
<i>Nepenthes</i>	Phytosanitary certificate not required.
<i>Nerium</i>	No additional declarations required.
<i>Nepeta</i>	Phytosanitary certificate not required.
<i>Nertera</i>	Phytosanitary certificate not required.
<i>Newtonia</i>	Phytosanitary certificate not required.
<i>Nierembergia</i>	Phytosanitary certificate not required.
<i>Nigella</i>	Phytosanitary certificate not required.
<i>Nolana</i>	Phytosanitary certificate not required.
<i>Nuxia</i>	No additional declarations required.
<i>Nuytsia</i>	No additional declarations required.
<i>Nymphaea</i>	No additional declarations required.
<i>Nyssa</i>	No additional declarations required.
<i>Ocimum</i>	Phytosanitary certificate not required.
<i>Oenothera</i>	Phytosanitary certificate not required.
<i>Olearia</i>	Phytosanitary certificate not required.

<i>Oncoba</i>	No additional declarations required.
<i>Ongokea</i>	No additional declarations required.
<i>Onobrychis</i>	No additional declarations required.
<i>Operculicarya</i>	No additional declarations required.
<i>Ophioglossum</i>	No additional declarations required.
Orchidaceae	No additional declarations required.
<i>Origanum</i>	Phytosanitary certificate not required.
<i>Ornithopus</i>	Phytosanitary certificate not required.
<i>Osteospermum</i>	Phytosanitary certificate not required.
<i>Ostrya</i>	No additional declarations required.
<i>Osyrius</i>	No additional declarations required.
<i>Oxypetalum</i>	Phytosanitary certificate not required.
<i>Pachypodium</i>	Phytosanitary certificate not required.
<i>Paeonia</i>	Phytosanitary certificate not required.
<i>Parnassia</i>	Phytosanitary certificate not required.
<i>Parinari</i>	No additional declarations required.
<i>Parkinsonia</i>	No additional declarations required.
<i>Parthenium</i>	Phytosanitary certificate not required.
<i>Pastinaca</i>	Phytosanitary certificate not required.
<i>Paulownia</i>	No additional declarations required.
<i>Pavetta</i>	No additional declarations required.
<i>Peganum</i>	No additional declarations required.
<i>Peltiphyllum</i>	Phytosanitary certificate not required.
<i>Peltophorum</i>	No additional declarations required.
<i>Pentas</i>	No additional declarations required.
<i>Penstemon</i>	Phytosanitary certificate not required.
<i>Peperomia</i>	Phytosanitary certificate not required.
<i>Perilla</i>	Phytosanitary certificate not required.
<i>Perovskia</i>	No additional declarations required.
<i>Persoonia</i>	No additional declarations required.
<i>Petalostemon</i>	Phytosanitary certificate not required.
<i>Petasites</i>	Phytosanitary certificate not required.

<i>Petrophila</i>	No additional declarations required.
<i>Petroselinum</i>	Phytosanitary certificate not required.
<i>Petunia</i>	Phytosanitary certificate not required.
<i>Peucedanum</i>	Phytosanitary certificate not required.
<i>Phacelia</i>	Phytosanitary certificate not required.
<i>Phalaris</i>	No additional declarations required.
<i>Philodendron</i>	Phytosanitary certificate not required.
<i>Phleum</i>	No additional declarations required.
<i>Phlomis</i>	No additional declarations required.
<i>Phlox</i>	No additional declarations required.
<i>Phormium</i>	No additional declarations required.
<i>Phygellus</i>	No additional declarations required.
<i>Phylloglossum</i>	Phytosanitary certificate not required.
<i>Phyllanthus</i>	No additional declarations required.
<i>Phyllocladus</i>	No additional declarations required.
<i>Physostegia</i>	Phytosanitary certificate not required.
<i>Phyteuma</i>	Phytosanitary certificate not required.
<i>Phytolacca</i>	No additional declarations required.
<i>Picea</i>	No additional declarations required.
<i>Pieris</i>	No additional declarations required.
<i>Pimpinella</i>	Phytosanitary certificate not required.
<i>Pinguicula</i>	Phytosanitary certificate not required.
<i>Piper</i>	No additional declarations required.
<i>Pisonia</i>	No additional declarations required.
<i>Pithecellobium</i>	No additional declarations required.
<i>Pittosporum</i>	No additional declarations required.
<i>Plagianthus</i>	No additional declarations required.
<i>Platycodon</i>	Phytosanitary certificate not required.
<i>Plectranthus</i>	No additional declarations required.
<i>Plumbago</i>	No additional declarations required.
<i>Plumeria</i>	No additional declarations required.
<i>Podocarpus</i>	No additional declarations required.

<i>Polemonium</i>	Phytosanitary certificate not required.
<i>Polyscias</i>	No additional declarations required.
<i>Pouteria</i>	No additional declarations required.
<i>Pothos</i>	Phytosanitary certificate not required.
<i>Pratia</i>	Phytosanitary certificate not required.
<i>Priurella</i>	No additional declarations required.
<i>Primula</i>	Phytosanitary certificate not required.
<i>Prunella</i>	Phytosanitary certificate not required.
<i>Prumnopitys</i>	No additional declarations required.
<i>Pseudolachnostylis</i>	Phytosanitary certificate not required.
<i>Pseudolarix</i>	No additional declarations required.
<i>Pseudotsuga</i>	No additional declarations required.
<i>Psilotum</i>	Phytosanitary certificate not required.
<i>Psiloxylon</i>	No additional declarations required.
<i>Psophocarpus</i>	No additional declarations required.
<i>Psoralea</i>	Phytosanitary certificate not required.
<i>Pterocarpus</i>	No additional declarations required.
<i>Pterodiscus</i>	No additional declarations required.
<i>Puccinellia</i>	No additional declarations required.
<i>Puelia</i>	No additional declarations required.
<i>Pulicaria</i>	Phytosanitary certificate not required.
<i>Pulmonaria</i>	Phytosanitary certificate not required.
<i>Pultenaea</i>	No additional declarations required.
<i>Punica</i>	No additional declarations required.
<i>Pyrostegia</i>	No additional declarations required.
<i>Radermachera</i>	No additional declarations required.
<i>Raphionacme</i>	No additional declarations required.
<i>Ravenala</i>	No additional declarations required.
<i>Rehmannia</i>	No additional declarations required.
<i>Reseda</i>	Phytosanitary certificate not required.
<i>Restio</i>	No additional declarations required.
<i>Reyesia</i>	Phytosanitary certificate not required.

<i>Rhamnus</i>	No additional declarations required.
<i>Rhodochiton</i>	No additional declarations required.
<i>Rhododendron</i>	No additional declarations required.
<i>Rhodohypoxis</i>	Phytosanitary certificate not required.
<i>Rhodosphaera</i>	No additional declarations required.
<i>Rhoeo</i>	Phytosanitary certificate not required.
<i>Rhytidocaulon</i>	No additional declarations required.
<i>Rogersia</i>	Phytosanitary certificate not required.
<i>Romneya</i>	Phytosanitary certificate not required.
<i>Roscoea</i>	No additional declarations required.
<i>Rosmarinus</i>	No additional declarations required.
<i>Rothmannia</i>	No additional declarations required.
<i>Rubia</i>	Phytosanitary certificate not required.
<i>Rudbeckia</i>	No additional declarations required.
<i>Ruellia</i>	Phytosanitary certificate not required.
<i>Ryania</i>	Phytosanitary certificate not required.
<i>Sagina</i>	Phytosanitary certificate not required.
<i>Salix</i>	No additional declarations required.
<i>Salpiglossus</i>	Phytosanitary certificate not required.
<i>Salvia</i>	Phytosanitary certificate not required.
<i>Samanea</i>	No additional declarations required.
<i>Sandoricum</i>	No additional declarations required.
<i>Santalum</i>	No additional declarations required.
<i>Sanvitalia</i>	Phytosanitary certificate not required.
<i>Saponaria</i>	Phytosanitary certificate not required.
<i>Sapium</i>	No additional declarations required.
<i>Saraca</i>	No additional declarations required.
<i>Sarcocephalus</i>	No additional declarations required.
<i>Sarcococca</i>	No additional declarations required.
<i>Sarracenia</i>	Phytosanitary certificate not required.
<i>Satureja</i>	Phytosanitary certificate not required.
<i>Saxifraga</i>	Phytosanitary certificate not required.

<i>Scabiosa</i>	Phytosanitary certificate not required.
<i>Scaevola</i>	No additional declarations required.
<i>Schefflera</i>	No additional declarations required.
<i>Schinus</i>	No additional declarations required.
<i>Schizaea</i>	Phytosanitary certificate not required.
<i>Schizanthus</i>	Phytosanitary certificate not required.
<i>Schizolobium</i>	No additional declarations required.
<i>Scindapsus</i>	Phytosanitary certificate not required.
<i>Scorzomera</i>	Phytosanitary certificate not required.
<i>Schoenocaulon</i>	No additional declarations required.
<i>Schotia</i>	No additional declarations required.
<i>Securidaca</i>	No additional declarations required.
<i>Sempervivum</i>	Phytosanitary certificate not required.
<i>Sesamum</i>	No additional declarations required.
<i>Shasta</i>	Phytosanitary certificate not required.
<i>Shepherdia</i>	No additional declarations required.
<i>Sicana</i>	No additional declarations required.
<i>Sidalcea</i>	Phytosanitary certificate not required.
<i>Silene</i>	Phytosanitary certificate not required.
<i>Simmondsia</i>	No additional declarations required.
<i>Sinapis</i>	No additional declarations required.
<i>Soleirolia</i>	No additional declarations required.
<i>Solidago</i>	Phytosanitary certificate not required.
<i>Solidaster</i>	Phytosanitary certificate not required.
<i>Solenostemon</i>	Phytosanitary certificate not required.
<i>Sollya</i>	No additional declarations required.
<i>Sophora</i>	No additional declarations required.
<i>Sparmannia</i>	No additional declarations required.
<i>Spartium</i>	No additional declarations required.
<i>Spathiphyllum</i>	No additional declarations required.
<i>Spiraea</i>	No additional declarations required.
<i>Spondias</i>	No additional declarations required.

<i>Spyridium</i>	No additional declarations required.
<i>Stachys</i>	Phytosanitary certificate not required.
<i>Stenocarpus</i>	No additional declarations required.
<i>Stephanotis</i>	No additional declarations required.
<i>Stereospermum</i>	No additional declarations required.
<i>Stevia</i>	Phytosanitary certificate not required.
<i>Stizolobium</i>	Phytosanitary certificate not required.
<i>Stokesia</i>	Phytosanitary certificate not required.
<i>Streptocarpus</i>	Phytosanitary certificate not required.
<i>Strychnos</i>	No additional declarations required.
<i>Stylosanthes</i>	No additional declarations required.
<i>Styphelia</i>	No additional declarations required.
<i>Swainsona</i>	No additional declarations required.
<i>Swartzia</i>	No additional declarations required.
<i>Swientenia</i>	No additional declarations required.
<i>Symphoricarpos</i>	No additional declarations required.
<i>Symphytum</i>	Phytosanitary certificate not required.
<i>Syngonium</i>	Phytosanitary certificate not required.
<i>Synsepalum</i>	No additional declarations required.
<i>Syzygium</i>	No additional declarations required.
<i>Tabebuia</i>	No additional declarations required.
<i>Tabernaemontana</i>	No additional declarations required.
<i>Tacca</i>	Phytosanitary certificate not required.
<i>Talinum</i>	Phytosanitary certificate not required.
<i>Tamarindus</i>	No additional declarations required.
<i>Tamarix</i>	No additional declarations required.
<i>Tanacetum</i>	Phytosanitary certificate not required.
Taxodiaceae	No additional declarations required.
<i>Telopea</i>	No additional declarations required.
<i>Templetonia</i>	No additional declarations required.
<i>Tephrosia</i>	No additional declarations required.
<i>Tetragonia</i>	Phytosanitary certificate not required.

<i>Tetraclinis</i>	No additional declarations required.
<i>Tetrataxis</i>	No additional declarations required.
<i>Teucrium</i>	Phytosanitary certificate not required.
<i>Thalictrum</i>	Phytosanitary certificate not required.
<i>Thermopsis</i>	Phytosanitary certificate not required.
<i>Thespesia</i>	No additional declarations required.
<i>Thunbergia</i>	No additional declarations required.
<i>Thymophylla</i>	Phytosanitary certificate not required.
<i>Thymus</i>	Phytosanitary certificate not required.
<i>Tiarella</i>	Phytosanitary certificate not required.
<i>Tibouchina</i>	Phytosanitary certificate not required.
<i>Tilia</i>	No additional declarations required.
<i>Tmesipterus</i>	No additional declarations required.
<i>Torenia</i>	Phytosanitary certificate not required.
<i>Trachelium</i>	Phytosanitary certificate not required.
<i>Trachymene</i>	Phytosanitary certificate not required.
<i>Tradescantia</i>	Phytosanitary certificate not required.
<i>Tragopogon</i>	Phytosanitary certificate not required.
<i>Triaspis</i>	No additional declarations required.
<i>Trigonella</i>	Phytosanitary certificate not required.
<i>Tristania</i>	No additional declarations required.
<i>Tritonia</i>	Phytosanitary certificate not required.
<i>Tropaeolum</i>	Phytosanitary certificate not required.
<i>Tsuga</i>	No additional declarations required.
<i>Turraea</i>	No additional declarations required.
<i>Umbellularia</i>	No additional declarations required.
<i>Uncarina</i>	No additional declarations required.
<i>Ursinia</i>	Phytosanitary certificate not required.
<i>Utricularia</i>	Phytosanitary certificate not required.
<i>Valeriana</i>	No additional declarations required.
<i>Valerianella</i>	Phytosanitary certificate not required.
<i>Vangueria</i>	No additional declarations required.

<i>Verbascum</i>	No additional declarations required.
<i>Verbena</i>	Phytosanitary certificate not required.
<i>Vernonia</i>	No additional declarations required.
<i>Veronica</i>	Phytosanitary certificate not required.
<i>Verticordia</i>	No additional declarations required.
<i>Viburnum</i>	No additional declarations required.
<i>Victoria</i>	No additional declarations required.
<i>Villarisa</i>	Phytosanitary certificate not required.
<i>Vinca</i>	Phytosanitary certificate not required.
<i>Viscaria</i>	Phytosanitary certificate not required.
<i>Vitex</i>	No additional declarations required.
<i>Voandzeia</i>	Phytosanitary certificate not required.
<i>Wahlenbergia</i>	Phytosanitary certificate not required.
<i>Wasabia</i>	Phytosanitary certificate not required.
<i>Warburgia</i>	No additional declarations required.
<i>Weinmannia</i>	No additional declarations required.
<i>Welwitschia</i>	No additional declarations required.
<i>Wisteria</i>	No additional declarations required.
<i>Witheringia</i>	No additional declarations required.
<i>Xanthisma</i>	Phytosanitary certificate not required.
<i>Xeranthemum</i>	Phytosanitary certificate not required.
<i>Zinnia</i>	Phytosanitary certificate not required.

4.6 Herbs, Spices, Medicinal Plants, & Plants for Animal & Human Consumption (Including Seed for Consumption)

4.6.1 Herbs, Spices, Medicinal Plants, & Plants for Animal & Human Consumption (Including Seed for Consumption)

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Herbs, Spices, Medicinal Plants, and Plants for Animal & Human Consumption (Including Seed for Consumption) require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate.

<i>Acanthus mollis</i>	No additional declarations required.
<i>Achillea</i>	No additional declarations required.
<i>Aconitum</i>	No additional declarations required.
<i>Actaea spicata</i>	No additional declarations required.
<i>Agar</i>	No additional declarations required.
<i>Agastache foeniculum</i>	No additional declarations required.
<i>Agrimonia eupatoria</i>	No additional declarations required.
<i>Ajuga</i>	No additional declarations required.
<i>Alcea rosea</i>	No additional declarations required.
<i>Alchemilla vulgaris</i>	No additional declarations required.
<i>Allium</i> (dried, flaked, powdered)	No additional declarations required.
<i>Aloe</i>	No additional declarations required.
<i>Althaea</i>	No additional declarations required.
<i>Anacardium</i> (nuts only)	No additional declarations required.
<i>Anagallis arvensis</i>	No additional declarations required.
<i>Anchusa</i>	No additional declarations required.
<i>Androstephium violaceum</i>	No additional declarations required.
<i>Anemone</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Anethum</i>	No additional declarations required.
<i>Angelica</i>	No additional declarations required.
<i>Antennaria dioica</i>	No additional declarations required.

<i>Anthemis tinctoria</i>	No additional declarations required.
<i>Anthriscus</i>	No additional declarations required.
<i>Anthyllis</i>	No additional declarations required.
<i>Apium graveolens</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Aquilegia vulgaris</i>	No additional declarations required.
<i>Arabis</i>	No additional declarations required.
<i>Aralia</i>	No additional declarations required.
<i>Arctotis acaulis</i>	No additional declarations required.
<i>Areca catechu</i> (nuts only)	No additional declarations required.
<i>Armeria</i>	No additional declarations required.
<i>Arnica montana</i>	No additional declarations required.
<i>Asarum canadense</i>	No additional declarations required.
<i>Asclepias tuberosa</i>	No additional declarations required.
<i>Asperula</i>	No additional declarations required.
<i>Astragalus</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Atropa</i>	No additional declarations required.
<i>Baptisia</i>	No additional declarations required.
<i>Barbarea verna</i>	No additional declarations required.
<i>Bellis perennis</i>	No additional declarations required.
<i>Borago officinalis</i>	No additional declarations required.
<i>Brassica</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Bryonia dioica</i>	No additional declarations required.
<i>Cajanus</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Calendula</i>	No additional declarations required.
<i>Campanula</i>	No additional declarations required.
<i>Canavalia</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Capparis spinosa</i>	No additional declarations required.

<i>Carthamus tinctorius</i>	No additional declarations required.
<i>Carum carvi</i>	No additional declarations required.
<i>Carya</i> (nuts only)	No additional declarations required.
<i>Castanea</i>	No additional declarations required.
<i>Catharanus roseus</i>	No additional declarations required.
<i>Ceanothus americanus</i>	No additional declarations required.
<i>Cedronella tryphylla</i>	No additional declarations required.
<i>Centaurea cyanus</i>	No additional declarations required.
<i>Centaureum erythraea</i>	No additional declarations required.
<i>Ceratonia siliqua</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cheiranthus cheiri</i>	No additional declarations required.
<i>Chelidonium</i>	No additional declarations required.
<i>Chenopodium</i>	No additional declarations required.
<i>Chrysanthemum</i>	No additional declarations required.
<i>Cicer</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cichorium intybus</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Cimicifuga racemosa</i>	No additional declarations required.
<i>Cinnamomum</i>	No additional declarations required.
<i>Cnicus benedictus</i>	No additional declarations required.
<i>Cocos nucifera</i> (nuts only)	No additional declarations required.
<i>Coffea</i> (Roasted beans only)	No additional declarations required.
<i>Colchicum autumnale</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Collinsonia canadensis</i>	No additional declarations required.
<i>Colutea</i>	No additional declarations required.
<i>Conium maculatum</i>	No additional declarations required.
<i>Convallaria</i>	No additional declarations required.
<i>Coreopsis tinctoria</i>	No additional declarations required.
<i>Coriandrum sativum</i>	No additional declarations required.

<i>Cornus florida</i>	No additional declarations required.
<i>Corylus</i> (nuts only)	No additional declarations required.
<i>Crithmum maritimum</i>	No additional declarations required.
<i>Crocus sativus</i>	No additional declarations required.
<i>Croton</i>	No additional declarations required.
<i>Cryptotaenia japonica</i>	No additional declarations required.
Cucurbitaceae (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cuminum cyminum</i>	No additional declarations required.
<i>Cupressus</i>	No additional declarations required.
<i>Curcuma longa</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Cyamopsis</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Cynara</i>	No additional declarations required.
<i>Danae racemosa</i>	No additional declarations required.
<i>Delphinium grandiflorum</i>	No additional declarations required.
<i>Dianthus</i>	No additional declarations required.
<i>Dictamnus albus</i>	No additional declarations required.
<i>Digitalis</i>	No additional declarations required.
<i>Dipsacus</i> (flower head)	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with Methyl bromide gas. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Dolichos</i>	No additional declarations required.
<i>Dracocephalum moldavica</i>	No additional declarations required.
<i>Elettaria cardemomum</i>	No additional declarations required.
<i>Epilobium</i>	No additional declarations required.
<i>Erica</i>	No additional declarations required.
<i>Erigeron</i>	No additional declarations required.
<i>Eugenia caryophyllata</i>	No additional declarations required.
<i>Euphorbia</i>	No additional declarations required.

<i>Fagopyrum</i>	No additional declarations required.
<i>Filipendula</i>	No additional declarations required.
<i>Foeniculum vulgare</i>	No additional declarations required.
<i>Galega officinale</i>	No additional declarations required.
<i>Galium</i>	No additional declarations required.
<i>Genista tinctoria</i>	No additional declarations required.
<i>Gentiana</i>	No additional declarations required.
<i>Geranium</i>	No additional declarations required.
<i>Glycines</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Glycyrrhiza glabra</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Gnaphalium</i>	No additional declarations required.
<i>Gossypium</i>	No additional declarations required. T2 = consignment was treated with a wide spectrum insecticide or fumigant. D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.
<i>Gratiola officinalis</i>	No additional declarations required.
<i>Grindelia robusta</i>	No additional declarations required.
<i>Guizotia abyssinica</i>	No additional declarations required.
<i>Hamamelis virginiana</i>	No additional declarations required.
<i>Helianthemum</i>	No additional declarations required.
<i>Heliotropium</i>	No additional declarations required.
<i>Helleborus niger</i>	No additional declarations required.
<i>Herniaria</i>	No additional declarations required.
<i>Heuchera</i>	No additional declarations required.
<i>Hibiscus sabdariffa</i>	No additional declarations required.
<i>Hieracium</i>	No additional declarations required.
<i>Hordeum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Humulus</i>	No additional declarations required.
<i>Hydrocotyle asiatica</i>	No additional declarations required.
<i>Hyoscyamus niger</i>	No additional declarations required.

<i>Hypericum</i>	No additional declarations required.
<i>Hyssopus officinalis</i>	No additional declarations required.
<i>Iberis umbellata</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Ilex paraguariensis</i>	No additional declarations required.
<i>Illicium verum</i>	No additional declarations required.
<i>Indigofera</i>	No additional declarations required.
<i>Inula helenium</i>	No additional declarations required.
<i>Iris</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Isatis tinctoria</i>	No additional declarations required.
<i>Jasminum</i>	No additional declarations required.
<i>Juglans</i> (nuts only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Juniperus communis</i>	No additional declarations required.
<i>Lamium</i>	No additional declarations required.
<i>Laurus</i>	No additional declarations required.
<i>Lavandula</i>	No additional declarations required.
<i>Lawsonia inermis</i>	No additional declarations required.
<i>Lens</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Leontopodium alpinum</i>	No additional declarations required.
<i>Leonurus</i>	No additional declarations required.
<i>Lepidium sativum</i>	No additional declarations required.
<i>Levisticum officinale</i>	No additional declarations required.
<i>Liatris</i>	No additional declarations required.
<i>Ligustrum</i>	No additional declarations required.
<i>Linaria</i>	No additional declarations required.
<i>Linum usitatissimum</i>	No additional declarations required.
<i>Lobelia</i>	No additional declarations required.
<i>Lophocarpus</i>	No additional declarations required.
<i>Lotus</i>	No additional declarations required.

<i>Lupinus</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Lychnis chalcedonica</i>	No additional declarations required.
<i>Lycium chinense</i>	No additional declarations required.
<i>Lycopus europaeus</i>	No additional declarations required.
<i>Lysimachia nummularia</i>	No additional declarations required.
<i>Lythrum salicaria</i>	No additional declarations required.
<i>Macadamia</i> (nuts only)	No additional declarations required.
<i>Maclura pomifera</i>	No additional declarations required.
<i>Mangifera</i> (seed only)	No additional declarations required.
<i>Malva</i>	No additional declarations required.
<i>Manihot esculenta</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Maranta arundinacea</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Marrubium vulgare</i>	No additional declarations required.
<i>Matricaria chamomilla</i>	No additional declarations required.
<i>Melilotus officinalis</i>	No additional declarations required.
<i>Melissa officinalis</i>	No additional declarations required.
<i>Mentha</i>	No additional declarations required.
<i>Metroxylon rumphii</i>	No additional declarations required.
<i>Mirabilis</i>	No additional declarations required.
<i>Molucella laevis</i>	No additional declarations required.
<i>Monarda didyma</i>	No additional declarations required.
<i>Morus nigra</i> (dried)	No additional declarations required.
<i>Myrica pennsylvanica</i>	No additional declarations required.
<i>Myristica fragrans</i>	No additional declarations required.
<i>Myrrhis odorata</i>	No additional declarations required.
<i>Myrtus communis</i>	No additional declarations required.
<i>Nasturtium officinale</i>	No additional declarations required.
<i>Nepenthes</i>	No additional declarations required.
<i>Nepeta</i>	No additional declarations required.
<i>Nigella damascena</i>	No additional declarations required.

<i>Ocimum</i>	No additional declarations required.
<i>Ononis</i>	No additional declarations required.
<i>Origanum</i>	No additional declarations required.
<i>Oryza</i> (grain only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Panax</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Papaver somniferum</i>	No additional declarations required. T7 = consignment concerned was treated by an appropriate fumigation with MBr gas. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Paris quadrifolia</i>	No additional declarations required.
<i>Pelargonium</i>	No additional declarations required.
<i>Perilla frutescens</i>	No additional declarations required.
<i>Petroselinum crispum</i>	No additional declarations required.
<i>Peucedanum</i>	No additional declarations required.
<i>Phalaris canariensis</i>	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Phaseolus</i>	No additional declarations required.
<i>Piper</i>	No additional declarations required.
<i>Pisum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Ricinus</i>	No additional declarations required.
<i>Saxifraga</i>	No additional declarations required.
<i>Secale</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Sedum</i>	No additional declarations required.
<i>Sempervivum</i>	No additional declarations required.
<i>Sesamum</i>	No additional declarations required.
<i>Silybum marianum</i>	No additional declarations required.
<i>Smilax</i>	No additional declarations required.
<i>Solidago virgaurea</i>	No additional declarations required.

<i>Sorghum</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Stachys</i>	AD11:- The consignment is free from soil or growth mediums.
<i>Symphytum</i>	No additional declarations required.
<i>Tagetes</i>	No additional declarations required.
<i>Taraxacum officinale</i>	No additional declarations required.
<i>Teucrium</i>	No additional declarations required.
<i>Tetragonia tetragonioides</i>	No additional declarations required.
<i>Theobroma</i>	No additional declarations required.
<i>Thymus</i>	No additional declarations required.
<i>Tilia cordata</i>	No additional declarations required.
<i>Trigonella foenum-graecum</i>	No additional declarations required.
<i>Tropaeolum</i>	No additional declarations required.
<i>Tussilago farfara</i>	No additional declarations required.
<i>Urginea maritima</i>	No additional declarations required.
<i>Urtica</i>	No additional declarations required.
<i>Valeriana</i>	No additional declarations required.
<i>Valerianella locusta</i>	No additional declarations required.
<i>Vanilla</i>	No additional declarations required.
<i>Verbascum</i>	No additional declarations required.
<i>Verbena</i>	No additional declarations required.
<i>Vernonia</i>	No additional declarations required.
<i>Veronica</i>	No additional declarations required.
<i>Viburnum opulus</i>	No additional declarations required.
<i>Vigna</i> (seed only)	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)
<i>Vinca</i>	No additional declarations required.
<i>Viola</i>	No additional declarations required.
<i>Voandzeia</i>	No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)

Yucca glauca

No additional declarations required. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)

Zea

AD2:- New Zealand is free from *Cephalosporium maydis*, and *Peronosclerospora maydis*. AD3:- The area of production is free from *Erwinia stewartii*. AD4:- The consignment is free from *Prostephanus truncatus*. (D = the importer of controlled goods is required to submit a declaration on a form obtained from the executive officer which declares the consignment is for immediate re-export or for purposes other than cultivation.)

4.7 Wood, and Unmanufactured Wood Products

4.7.1 Wood and Unmanufactured Wood products

Import permits are not required for the commodities listed below. All other commodities in the commodity sub-class Wood and Unmanufactured Wood products require import permits (refer to sections 2.3, 2.4, 2.5, 2.6, and 2.7). Phytosanitary certificate required, additional declarations (AD's) and/or treatments as specified below. Treatments are not required to be written on the phytosanitary certificates as AD's if "No additional declarations required" is stated. Treatment details (product, rate/dose etc) are to be specified in the treatment section of the phytosanitary certificate. **All wood and unmanufactured wood products must be free from bark.**

<i>Abies</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4:- The consignment is free from <i>Gremmeniella abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Acacia</i>	No additional declarations required.
<i>Acer</i>	AD3:- The area of production is free from <i>Ceratocystis coerulescens</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Azelia</i>	No additional declarations required.
<i>Agathis</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Alectryon</i>	No additional declarations required.
<i>Amburana</i>	No additional declarations required.
<i>Anisoptera</i>	No additional declarations required.
<i>Anthocleista</i>	No additional declarations required.
<i>Araucaria</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Artocarpus</i>	No additional declarations required.
<i>Aucoumea</i>	No additional declarations required.
<i>Baeckea</i>	No additional declarations required.
<i>Baikrera</i>	No additional declarations required.
<i>Balfourodendron</i>	No additional declarations required.
<i>Berchemia</i>	No additional declarations required.
<i>Betula</i>	No additional declarations required.
<i>Bowdichia</i>	No additional declarations required.
<i>Brachystegia</i>	No additional declarations required.
<i>Buxaceae</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.

<i>Callitris</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Cariniana</i>	No additional declarations required.
<i>Carya</i>	No additional declarations required.
<i>Castanea</i>	AD3:- The area of production is free from <i>Ceratocystis fagacearum</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Castanopsis</i>	AD3:- The area of production is free from <i>Ceratocystis fagacearum</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Casuarina</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Cedrus</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4:- The consignment is free from <i>Gremmeniella abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Celtis</i>	AD3:- The area of production is free from <i>Ceratocystis ulmi</i> . AD4:- The consignment is free from <i>Lymantria disparina</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Copaifera</i>	No additional declarations required.
<i>Chlorophora</i>	No additional declarations required.
<i>Cupressus</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Cuphea</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Dipteryx</i>	No additional declarations required.
<i>Dyera</i>	No additional declarations required.
<i>Entandophragma</i>	No additional declarations required.
<i>Eucalyptus</i>	No additional declarations required.
<i>Euclea</i>	No additional declarations required.
<i>Fagus</i>	AD3:- The area of production is free from <i>Ceratocystis ulmi</i> . AD4:- The consignment is free from <i>Lymantria disparina</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Fraxinus</i>	AD3:- The area of production is free from <i>Ceratocystis fagacearum</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Gonystylus</i>	No additional declarations required.
<i>Guibourtia</i>	No additional declarations required.
<i>Intsia</i>	No additional declarations required.
<i>Juglans</i>	No additional declarations required.

<i>Kalopanax</i>	No additional declarations required.
<i>Khaya</i>	No additional declarations required.
<i>Koompassia</i>	No additional declarations required.
<i>Larix</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4 The consignment is free from <i>Gremmeniela abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Lithocarpus</i>	AD3:- The area of production is free from <i>Ceratocystis fagacearum</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Millettia</i>	No additional declarations required.
<i>Perecopsis</i>	No additional declarations required.
<i>Phoebe</i>	No additional declarations required.
<i>Physocolumma</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Picea</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4 The consignment is free from <i>Gremmeniela abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Pinus</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4 The consignment is free from <i>Gremmeniela abeitina</i> & <i>Lymantria disparina</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Platanus</i>	No additional declarations required.
<i>Podocarpus</i>	No additional declarations required.
<i>Populus</i>	AD4:- The consignment is free from <i>Campanotus herculeanus</i>
<i>Pseudotsuga</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4 The consignment is free from <i>Gremmeniela abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Pterocarpus</i>	No additional declarations required.
<i>Pterygota</i>	No additional declarations required.
<i>Quercus</i>	AD3:- The area of production is free from <i>Ceratocystis fagacearum</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Rosaceae</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Rutaceae</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Sassafras</i>	No additional declarations required.
<i>Shorea</i>	No additional declarations required.

<i>Spirostachys</i>	No additional declarations required.
<i>Sweetenia</i>	No additional declarations required.
<i>Tarrieta</i>	No additional declarations required.
<i>Taxaceae</i>	No additional declarations required.
<i>Taxodiaceae</i>	No additional declarations required.
<i>Tectona</i>	No additional declarations required.
<i>Terminalia</i>	AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Theobroma</i>	AD3:-The area of production is free from <i>Ceratocystis fimbriata</i> or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Thuja</i>	No additional declarations required.
<i>Tieghemella</i>	No additional declarations required.
<i>Tilia</i>	No additional declarations required.
<i>Triplochiton</i>	No additional declarations required.
<i>Tsuga</i>	AD3:- The area of production is free from <i>Bursaphelenchus xylophilus</i> , & <i>Sirex noctilio</i> . AD4:- The consignment is free from <i>Gremmeniella abeitina</i> & <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Ulmus</i>	AD3:- The area of production is free from <i>Ceratocystis ulmi</i> . AD4:- The consignment is free from <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.
<i>Umbellularia</i>	No additional declarations required.
<i>Zelkova</i>	AD3:- The area of production is free from <i>Ceratocystis ulmi</i> . AD4:- The consignment is free from <i>Lymantria disparina</i> , or AD9:- The timber comprising this consignment was kiln treated to moisture content of < 20%.